

OSPINA SUNIOR'S

PRODUCTS

CJ WW

# BUSOGA REGION JOINT EXAMINATION BOARD

Uganda Advanced Certificate of Education MOCK EXAMINATION 2017

**PHYSICS** 

Paper 1

2 hours 30 minutes

MONDAY: 14<sup>TH</sup>/08/2017

MORNING: 9:00AM - 11:30AM

# INSTRUCTIONS TO CANDIDATES

- Attempt any five questions, including at least one question from each of the sections A, B and C but NOT more than two questions from each section.
- Silent non-programmable scientific electronic calculators may be used.
- Where necessary assume:

| Where necessary                       |                      | 2  |
|---------------------------------------|----------------------|--|
| Acceleration due to gravity (g)       | to manife            | $9.81 ms^{-2}$                                     |
| Electronic mass $(M_e)$               | agentina<br>america  | $9.11 \times 10^{-31} kg$                          |
| One electron volt (ev)                | =                    | $1.6 \times 10^{-19} J$                            |
| Planck's constant (h)                 |                      | $6.63 \times 10^{-34} Js$                          |
| Radius of the earth *                 | money<br>u minut     | $6.4 \times 10^6 m$                                |
| Radius of the sum                     | =                    | $6.96 \times 10^8 m$                               |
| Speed of light in a vacuum (c)        | =                    | $3.0 \times 10^8 m$                                |
| Stefan's constant; \sigma             | =                    | $5.67 \times 10^{-8} W  m^{-2} K^{-4}$             |
| Charge to mass ratio $(\frac{e}{m})$  | ==                   | 1.8 x 10 <sup>-11</sup> Ckg <sup>-1</sup>          |
| Universal gravitational constant (G)  | =                    | $6.67 \times 10^{-11}  \text{Nm}^2 \text{kg}^{-2}$ |
| Avogadro's number $(\tilde{N}_A)$     | , , <del>=</del>     | $6.02 \times 10^{23} mol^{-1}$                     |
| lev                                   | Manifer<br>page 1941 | 931MeV   |
| . Molar gas constant (R)              | ===                  | $8.31  JK^{-1} mot^{-1}$                           |
| Specific heat capacity of copper      | 14/4mm               | $3.70 \times 10^2 J kg^{-1} k^{-1}$                |
|                                       | ===                  | $1.66 \times 10^{-27} kg$                          |
| Unified mass unit (  U)               | ===                  | 4200Jkg <sup>-1</sup> K <sup>-1</sup>              |
| Specific heat capacity of water       | =                    | $1.47 \times 10^{11} m$                            |
| Radius of earth's orbit about the sun |                      | 1.7/ 2.10 11                                       |

#### SECTION A:

- 1. a) Define the following terms as applied to oscillatory motion.
  - i. Amplitude (01mk)
  - ii. Period (01mk)
  - b) State four characteristics of simple harmonic motion (02mks)
  - c) A mass, M is suspended from a rigid support by a string of length, l. The mass is pulled a side so that the string makes an angle,  $\theta$  with the vertical and then released.
  - i) Show that the mass execute simple harmonic motion with a period,

$$T = 2\pi \sqrt{\frac{l}{g}}$$
 (05mks)

- ii) Explain why this mass comes to a stop after a short time. (02mks)
- d) A piston in a car engine performs a simple harmonic motion of frequency 12.5Hz. If the mass of the piston is 0.50kg and its amplitude of vibration is 45mm, find the maximum force on the piston (3mks)
- e) Describe an experiment to determine the acceleration due to gravity, g using a spiral spring of known force constant. (06mks)
- 2. a) Define viscosity of a fluid (01mk)
  - b) (i) Derive an expression for terminal velocity attained by a sphere of density  $\delta$ , and radius, a falling through a fluid of density  $\rho$  and coefficient of viscosity  $\eta$ .
  - ii) Explain the variation of viscosity of a liquid with temperature (02mks)
  - c) i) State the laws of solid friction (02mks)
  - ii) With the aid of a well labeled diagram, describe an experiment to determine the coefficient of kinetic function between two surfaces. (05mks)

|   | 1 sland inclined at 30° to the horizontal.                          | If the  |
|---|---|---------|
| d | A body slides down a rough plane inclined at 30° to the horizontal. | find    |
| / | A body slides down a rough pro-                                     | , 11119 |
|   | coefficient of killede meters as                                    |         |
|   | the velocity after it has travelled 6m along the plane.             | (0.1 mk |

3. a)Define gravitational potential at a point

(01mk)

- b) (i) Describe the energy changes that occur as a space-craft falls towards the earth well away from the earth's atmosphere.
  - ii) Calculate the minimum energy required to project a space craft of mass
  - $2.0 \times 10^6 kg$  from the surface of the earth so that it escapes completely from (04mks) the influence of the earth's gravitational field.

(Radius of the earth = 6400km, Mass of the earth =  $6.0 \times 10^{24}$ kg)

- State Newton's law of gravitation and give the units of the gravitational ì. (02mks) constant.
- Briefly describe an experiment to determine the universal gravitational ii. (06mks) constant.
- (02mks) Explain why the moon has no atmosphere iii.
- √4. a) i) Define Young's modulus and strain of an elastic material (02mks)
  - ii) Derive an expression for the energy stored per unit volume of a stretched (03mks) wire in terms of Young's modulus and strain.
  - b) Describe an experiment with the aid of a labeled diagram, to investigate the relationship between tensile stress and tensile strain of a steal wire
  - c) A steel wire in a guitar has a length of 0.75m and mass per unit length of  $8.0 \times 10^{-4} \text{kgm}^{-1}$ . Its tension is adjusted so that it vibrates at a fundamental frequency of 256Hz when plucked.
  - (03mks) i) Calculate the value of the tension in the wire
  - ii) Calculate the energy stored per unit volume in the stretched wire if it has (02mks)a cross-sectional area  $1.0 \times 10^{-7} \text{m}^2$
  - iii) The wire snaps at one of the bridges. Find the total energy stored in the wire and hence estimate the initial speed of the wire. (Young's modulus of (03mks) steel = 2.0 x 10 "Pa) Page 3

## SECTION B:

5. a) (i) Define thermal radiation

(01mk)

ii) Describe how infra-red radiations can be detected

(4mks)

b) i) Explain the mechanism of heat transfer in insulators

(03mks)

ii) Describe how the thermal conductivity of rubber can be determined

(07mks)

- a) A rain drop of mass m falls to the ground at its terminal velocity V. The specific heat capacity of water is C and the acceleration of free fall is g. Given that 25% of the energy is retained when it strikes the ground, derive an expression for the temperature rise of the rain drop.
- b) A well lagged copper bar having a uniform cross-sectional area of 2.0 x 10-4m<sup>2</sup> is heated at one end and is cooled by a flow of water at the other end. The temperature gradient along the bar is 2.1Kcm-1 when the rate of water flow is 1.0gs-1. If the heat taken up by the water raises its temperature by 4.0K, calculate the thermal conductivity of copper (03mks) (specific heat capacity of water is 4.2KJkg-1K-1.
- √ 6. a) (i) Define the term specific heat capacity, internal energy and state their (03mks) units.
  - (ii) Why is the distinction between specific heat capacity at constant pressure and that at constant volume important for gases, but less (04mks) important for solids and liquids?
  - b) Explain why the temperature of a liquid does not change when the liquid (02mks) is boiling?
  - c) One kilogram of water is converted to steam at a temperature of 100°C and a pressure of 1.0 x  $10^5$ Pa. If the density of steam is 0.58kgm<sup>-3</sup> and the specific latent heat of vaporization of water is 2.3 x 106Jkg-1. Calculate the
  - i) external work done

(04mks)

ii) internal energy

(03mks)

a) Explain why the specific latent heat of fusion and specific latent heat of vaporization of a substance at the same pressure are different. (04 mks)

- 7. a) (i) Distinguish between isothermal and adiabatic changes (02mks)
  - (ii) State the condition under which an adiabatic change can take place (02mks)
  - (iii) Explain why the passage of sound waves through air is considered as an adiabatic process. (03mks)
  - b) A fixed mass of an ideal gas whose ratio Cp: Cv is 5:3 has a temperature of  $27^{\circ}$ C, volume of  $6.4 \times 10^{-2} \text{m}^{3}$  and pressure  $243 \text{Nm}^{-2}$ . It undergoes a reversible adiabatic compression to a volume of  $2.7 \times 10^{-2} \text{m}^{3}$ . The gas is then expanded isothermally to the original volume.
  - i) Show an indicator diagram in the above process (02mks)
  - ii) Calculate the pressure at the end of the compression (03mks)
  - c) (i) What is meant by the thermometric property? (01mk)
    - (ii) List four thermometric properties which are used in thermometry (01mk)
      - iii) The resistance of a thermistor over a limited range of temperature is given by the equation;  $R = \frac{C}{T 203}$ , where C is a constant and T is the absolute temperature. T = 310 K. (05mks)

### SECTION C

8. a) (i) What is meant by mass defect?

(01mk)

- ii) Sketch a graph showing how binding energy per nucleon varies with mass number and explain its main features (03mks)
- iii) Find the binding energy per nucleon of  $\frac{56}{26}$  Fe given that

mass of 1 proton = 1.007825u

mass of 1 neutron = 1.008665u

(03mks)

[1u = 931 MeV]

b) (i) Show that when an alpha particle collides head - on with an atom of atomic number Z, the closest distance of approach to the nu

given by 
$$X_0 = \frac{Ze^2}{\prod \sum_0 mV^2}$$

e is electronic change Where:

 $\Sigma_0$  is permittivity of free space

m is mass of alpha particle

v is initial speed of the alpha particle

(4mks)

ii) In a head-on collision between an alpha particle and a gold nucleus, the minimum distance of approach is 5 x 10<sup>-14</sup>m. Calculate the energy of the alpha particle (in MeV).

(Atomic number of gold = 79).

(03mks)

c) With the aid of a diagram, explain how an ionization chamber works.

(06mks)

- 9. (a) Define the following terms as used in photoelectric emission
  - Threshold frequency

(01mk)

(01mk)

b) Describe an experiment you would use to determine plank's constant, h.

(06mks)

(a) Light of wave length  $6.0 \times 10^2$ nm and intensify  $1 \times 10^{-5}$ Wm<sup>-2</sup> falls on a photo-sensitive surface of area 2.0cm<sup>2</sup> and work function 1.9eV. If 80% of the photons can cause electron emission, find the;

Photon energy for this light

(02mks)

Photon current that results

(04mks)

i) Explain how the line spectra in an x-ray tube is produced

(03mks)

ii) Electrons in an x-ray tube are accelerated through a pd of 14kv.

Calculate the minimum wave length of the x-rays that can be produced.

(03mks)

|           |   | $-e^2$                             |
|-----------|---|------------------------------------|
| 10. a) Sł | how that the total energy of an electron in an orbit is given by  | $8\Pi\Sigma_0 r$                   |
| 10. 4     |   | (05mks)                            |
| b) The    | energy levels shown in figure 1 below are of an atom of a certa   | ain                                |
|           |   |                                    |
| n = 4     | nt X. O eV1 eV  |                                    |
| n = 3     | -1 eV<br>-3 eV<br>-10eV   |                                    |
| n = 2     | -10eV   | 1                                  |
| n = 1     | Fig 1   |                                    |
| į.        | What is the shortest wave length of radiation that can be em X?  What region of the electromagnetic spectrum do you find the  | (03mks)<br>e radiation             |
| ii.       | What region of the electronia   | (01mk)                             |
|           | emitted in (i) above?   | (01mk)                             |
| c) (i)    | What are positive rays?   | sitive rays.                       |
| (ii) I    | Describe an experiment to determine the specific charge of pos  | (05mks)                            |
| 1         | An oil drop of mass $3.25 \times 10^{-15}$ kg falls vertically with uniform between two vertical parallel plates which are 2.0cm apart. W 1000V is applied between the plates, the drop moves towards | velocity hen a p.d of the negative |
|           |   |                                    |
|           | The why the vertical components   | (01mk)                             |
|           | unchanged.  | (04mks)                            |
|           | ii. Calculate the charge on the drop.   | •                                  |
|           | 4   |                                    |
|           | END   | l                                  |