

JINJA JOINT EXAMINATION BOARD

CHEMISTRY PAPER 2

545/2

JULY/AUGUST - 2019

PROPOSSED MARKING GUIDE

	PROPOSSED MARKING GOIDE	
		MARKS
	SECTION A (50 MARKS)	1 mark
Nos.	(i) Ammonium chloride and sodium chloride or equivalent	1 mark
1.	/ (") 11/-4 and could or equivalent	1 mark
	(iii) Ethanol and water or petroleum or equivalent	1 mark
	Let N to the J Culphur of equivalent	1 mark
1	(iv) Iron and Sulphur of Equivalent (v) Sodium chloride and potassium chlorate or equivalent	5 marks
	(v) Southin emorate and participation of the control of the contro	1 mark
2 (2)	Heated magnesium, steam	
2. (a)	(i) $Mg(s) + H_2O(g)$ $\longrightarrow MgO(s) + H_2(g)$	2 1111111111111111111111111111111111111
(b)	Heated magnesium, steam (i) $Mg(s) + H_2O(g)$ (ii) $MgO(s) + H_2O(l)$ $\longrightarrow MgO(s) + H_2(g)$ $\longrightarrow Mg(OH)_2(aq)$	$1\frac{1}{2}$ marks $1\frac{1}{2}$ marks
	(ii) ingo(c) X	1 mark
(c)	Turns red litmus paper to blue	5 marks
(0)	X	
3. (a)	(i) The yield of ammonia decreases because the forward reaction is exothermic.	1 mark
J. (a)	(ii) The yield of ammonia increases because the forward reaction proceeds by decrease	1 mark
	in volume/ number of moles.	1
	(iii)The yield of ammonia increases because iron acts as a catalyst.	1 mark
(b)	(i) volume of ammonia = $\frac{2}{3}$ x 120 = 80 cm ³	
(0)	(i) volume of ammonia = $\frac{2}{3}$ x 120 = 80cm ³	l mark
	(ii) Volume of nitrogen = $\frac{1}{3}$ x 120 = 40 cm ³	1 mark
		5 marks
4. (a)	(i) F VX	$\frac{1}{2}$ mark
()	(ii) D	~
		$\frac{1}{2}$ mark
	(iii) A and C	1 mark
	(iv) B or E	1 -
	X X	$\frac{1}{2}$ mark
(b)	(i) FD ₂ , ionic or electrovalent	
	(ii) HD, covalent	1 mark
	X	1 mark
		$4\frac{1}{2}$ marks
. (a)	$CO_2(g) + Cs)$ $\rightarrow 2CO(g)$	- 4
. (2)	$CO_2(g) + Cs)$ $\longrightarrow 2CO(g)$	$1\frac{1}{2}$ marks
(b)	(i) To absorb carbon dioxide gas	1 1
(5)	(ii) The black solid turns to brown	$\frac{1}{2}$ mark
(c)	(1) A G	1 mark
(6)	· /	- mark
	moles $\frac{0.48}{64} = 0.0075$ $\frac{0.06}{16} = 0.00375$	1 .
	10	$\frac{1}{2}$ mark
	mole ratio $\frac{0.0075}{0.00375} = 2$ $\frac{0.00375}{0.00375} = 3$	-
	mole ratio $\frac{0.0075}{0.00375} = 2$ $\frac{0.00375}{0.00375} = 1$	$\frac{1}{2}$ mark
	· .	2 mark
	formula Cu ₂ O	
		1 mark
	(ii) Copper (I) oxide	12
	V V	$\frac{1}{2}$ mark
		5 mark

	$\frac{1}{2}$ mark
X CU(a) 1	1 mark
6. (a) (i) Ethanol CH ₂ = CH ₂ (g)	1 mark
(ii) CH ₂ CH ₂ OH(l) (b) (i) A compound which contains double bonds between the carbon atoms (b) A compound which contains double bonds between the carbon atoms (c) A cidified potassium permanganate or equivalent, it turns from purple to	$1\frac{1}{2}$ mark
(b) (i) A compound which contains double bonds between the carbon atoms (ii) Acidified potassium permanganate or equivalent, it turns from purple to	2
(b) (i) A compound when the co	1 ,
(II) Actorited positions (II)	$\frac{1}{2}$ mark
	$\frac{1}{2}$ mark
(c) (i) Polymerization (ii) marking polythene papers or equivalent.	2
(II) marking posture 1 1	1,
(iii) - It pollutes the environment, X	$\frac{1}{2}$ mark
(iii) - It pollutes the environment, v - It affects soil fertility or <i>equivalent</i> . Accept one correct answer	5 ¹ mark
	2 111411
7. (a) (i)	1
/ X ½ /	
* /x\ T \	1 mark
\ \ y \sigma	
X _X	1
(ii)	
/ X \	1
	1 mark
I IX N IV	1 mark
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	
\backslash \forall $/$ \checkmark	
(b) Oxygen is more reactive because little energy is required to break the double bond	
	1 mark
between the oxygen atoms.	
Nitrogen is unreactive because a lot / high energy is required to break the strong triple hand between the nitrogen atoms.	1 mark
(c) (i) Manufacture of ammonia by the Haber process	
(ii) Cutting and welding of metals, making of steel of equivalent	1 mark
(a) Iron (II) sulphide is a compound	5 marks
Iron and Sulphus is a compound	1 mark
Iron and Sulphur is a mixture or equivalent (b) (i) FeS(s) +H ₂ SO ₄ (aq) ————————————————————————————————————	$1\frac{1}{2}$ marks
120(V) + FEX (1.420) V	_
(ii) $Fc(s) + H_2SO_4(aq)$ FeSO ₄ (aq) + $H_2(g)$	$1\frac{1}{2}$ marks
	2
(c) (ii) It is poisonous	Imort
a) (i) Zinc carbonate	lmark
	5 marks
(ii) Zinc chloride	
$ZnCO_3(s) + 2HCl(aq)$ \longrightarrow $ZnCl_2(aq) + CO(cs) + IV O(s)$	1 mark
ZnCO ₃ (s) + 2HCl(aq) \longrightarrow ZnCl ₂ (aq) + CO ₂ (g) + H ₂ O(l) \checkmark Moles of carbon dioxide gas $\frac{501.76}{22,400} = 0.0224$ moles \checkmark Moles of Zinc carbonate that received $= 1.0024$	
Moles of Zing and sent at	lmark
Moles of Zinc carbonate that reacted = $1 \times 0.0224 = 0.0224$ moles	
Molecular mass - 577, GO	
Molecular mass of $ZnCO_3 = 65 + 12 + 48 = 125 gm$	1
Mass of ZnCO ₃ that reacted = $0.0224 \times 125 = 2.8 \text{gm}$	1 mark
2.0giii	
Mass of ZnCl ₂ in the mixture = $5.0 - 2.8 = 2.2$ gm	
5.0 - 2.0 = 1.70m A	1 mark
Percentage of ZNCl ₂ = $\frac{2.2}{2.2}$ v 102	
Percentage of $ZNCI_2 = \frac{2.2}{5.0} \times 100 = 44\%$	
Percentage of $ZNCl_2 = \frac{2.2}{5.0} \times 100 = 44\%$	
Percentage of ZNCI ₂ = $\frac{2.2}{5.0}$ x 100 = 44%	1 monte
Percentage of $ZNCl_2 = \frac{2.2}{5.0} \times 100 = 44\%$	1 mark 5 marks

	<i>V</i>	
10.	(a) Q is Oxygen gas	
	(a) Q is Oxygen gas V R is Hydrogen gas V (b) (i) $4OH'(aq)$ $O_2(g) + 2H_2O(1) + 4e$	l mark
'		2
	(ii) $2H^{+}(aq) + 2e$ Gas R – Haber process H ₂ (g)	$l^{\frac{1}{2}}$ mark
	(c) Gas R – Haber process	
į	- Hydrogenation	1/2 mark
	- Manufacture of hydrochloric acid	$\frac{1}{2}$ mark
	d) Platinum X	5 marks
	SECTION B (30 MARKS)	
11. (lmarks
	oxygen	
	(ii)	
	Thermometer	
į		-
ĺ	Thin tin can	
į		3 marks
	Water	
	ethanol flame	
İ		
i	· \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	1
	ethanol	
	working apparatus 1 mark	
	- A known mass (M gm) of water is added to a tin can and its initial tempt. Noted ti°C.	
	- the total mass of ethanol and the lamp is determined, W ₁ gm.	
	- the lamp is lit and whose heat is used to heat water, the water is stirred	
İ	- the maximum temperature of heated wateris measured and the flams put off - the mass of ethanol and lamp is again determined as W ₂ (g).	
ĺ	the mass of editation and famp is again determined as $W_2(g)$.	
į	RESULTS	
1	Mass of ethanol burnt = $W_1 - W_2 = W_{gm} \nearrow X$	6 marks
j	temperature cit. non $= t_2 - t_1 = \Delta $?	o marks
1	Specific heat capacity = C	
	Molecular mass of ethanol = W_R	
	Quantity of heat produced = $(M \times C \times \Delta T)$ Joules \checkmark	
	Wgm of ethanol produced (M x C x Δ T) Joules of heat = H W _R gm of ethanol will produce $H \times W_R$ Joules	1
ĺ	Wight of chianol with produce TX WR Joures	1
	Heat of combustion = - H x Wn Joules/ mole?	
Ì	Heat of combustion = $-\frac{H \times W_R}{W}$ Joules/ mole	
!		
(b)	Heat change = $200 \times 4.2 \times 30 = 25,200$ Joules	
	Molecular mass of ethanol = 46gm	
	• •	
	1120KJ of heat requires 46gm of ethanol	
	25.2KJ of heat will require 46 x 25.2 gm of ethanol	3 marks
	1120	
	= 1.035gm \(
(c)	- It provides information about energy content of food metalical	
	- It provides information about effectiveness of fuel.	2 marks
		1.5
		15 marks

	(a) In the solid state, the lead (II) and bromide ions are strongly attracted by strong	1 .
12.	In the solid state, the lead (II) and bromide ions are strongly	$3\frac{1}{2}$ marks
12.	(a) In the solid state, the lead (II) and bromide forms the solid state, the lead (II) and bromide forms are mobile. In the molten state, the heat breaks the force of attraction between the ions, they are free and mobile to conduct electricity.	
	and mobile to conduct electricity.	
	Water being a polar solvent causes hydrochloric to ionize forming hydrogen ions and	
(1	chloride ions in solution.	
	$V \longrightarrow W(aa) + CI(ag) \checkmark$	4 marks
	The leader on lone react with magnesium to form hydrogyn gas	4]]]a] K3
	Mg(s) + 2H ⁴ (aq) \longrightarrow Mg ²⁺ (aq) + H ₂ (g) \longrightarrow Mg(s) + 2H ⁴ (aq) \longrightarrow Mg(s) + 2H ⁴ (aq) \longrightarrow Mg(s) + Mg(s) + H ₂ (g) \longrightarrow Mg(s) + H ₂ (g) \longrightarrow Mg(s) + Mg	
	Methyl benzene is a non-polar solvent, it doesn't cause romand	
	gas, hence no hydrogen ions in solution.	
	The constitution and those institution calcium sulphate.	
(0	The reaction produces insoluble calcular surplicates $CaSO_4(s) + CO_2(g) + H_2O(1) \checkmark$	3 marks
	The reaction produces insoluble calcium sulphate, CaCO ₃ (s) + H ₂ SO ₄ (aq) — CaSO ₄ (s) + CO ₂ (g) + H ₂ O(l) This form a coating on calcium carbonate which prevents the sulphuric acid from getting	
	into contact with calcium carbonate, hence the reaction stops.	
		1 .
(d)	The white precipitate is due to the formation of insoluble zinc hydroxide,	$4\frac{1}{2}$ mark
	Zn^{2} (aq) + 2OH (aq) \longrightarrow $Zn(OH)_2(s)$ In excess ammonia, the zinc hydroxide dissolves forming a soluble complex which is	
	colourless	
		15 marks
13. (a	(i) A normal is a compound formed when all the ionizable hydrogen of an acid is	$l^{\frac{1}{2}}$ mark
	replaced by a metal or ammonium ions. Sodium sulphate or equivalent	
	(ii) A acid salt is a compound formed when part of the ionizable hydrogen of an acid is	$1\frac{1}{2}$ mark
	replaced by a metal. Sodium hydrogen sulphate or equivalent	_
	X X	
(b)	(i) To warm dilute nitric acid in a beaker is added lead (II) oxide and the mixture	
, ,	stirred. Lead (II) nitrate is formed by the reaction.	5 marks
	$PbO(s) +2HNO_3(aq)$ \longrightarrow $Pb(NO_3)_2(aq) + H_2O(l)$	5 marks
	Lead (II) oxide is added until in excess. The excess lead (II) oxide is filtered off, the filtrate is heated to saturation point, the	
	saturated solution is cooled to form crystals. The crystals filtered, washed & dried	.
	saturated solution is cooled to form crystais. The crystais interest, washed craited	
	(ii) – lead (II) carbonate	1 mark
	Lead (II) hydroxide	
(c)	(i) A white precipitate	1 mark
	(ii) Preparation of insoluble salts.	1 mark
(L)	25 1 225 1	
(d)	(i) Moles of sulphate ions = $\frac{25}{1000}$ x 1 = 0.025 moles	
	40 X	
	Moles of lead (II) ions = $\frac{40}{1000}$ x 0.1 = 0.02 moles $\sqrt{}$	$1\frac{1}{2}$ mark
	Mole ratio 1:1	2
	X	
	Moles of sulphate ions that reacted = 0.02 moles \checkmark	
	(i) M. I (1-1/10-11-1-6	
	(ii) Moles of lead (II) sulphate formed = 0.02 moles	$2\frac{1}{2}$ mark
	Molecular mass of lead (II) sulphate = $207 + 32 + 64 = 303 \text{gm}$	2 11101 K
	Profession mass of lead (11) sulphate $-207 + 32 + 64 = 303 \text{gm} \checkmark$	
	Mass of lead (II) sulphate = $0.02 \times 303 = 6.02 \text{ gm}$	
	Contract of the second of the	15 marks
		15 marks

	X	
14. (a)	(i) Iron (II) carbonate, FeCO ₃	1 mark
	(ii) To convert Iron (II) carbonate to Iron (II) oxide	l mark
(b)	(i) Coke burns in hot air to form carbon dioxide gas $C(s) + O_2(g) \longrightarrow CO_2(g)$ Carbon dioxide gas is reduced by hot coke to carbon monoxide. $2CO_2(g) + C(s) \longrightarrow 2CO(g)$ Carbon monoxide reduces the hot iron ore to iron, $Fe_2O_3(s) + 3CO(g) \longrightarrow 2Fe(l) + 3CO_2(g)$	6 marks
(c)	(ii) Heated limestone decomposes to form calcium oxide and carbon dioxide gas. CaCO ₃ (s)	4 marks
	(ii) It does not assitute of	1 mark
1.1 Character	(ii) It does not easily rust	
St. C. St. Co. Co. Co. Co. Co. Co. Co. Co. Co. Co	It has high strength	
or above a		
		15 mark

END