Name:	
Signature	School: ,
Nigitates viiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiiii	

545/2 CHEMISTRY Paper 2 Jul / Aug. 2022 2 hours



### UGANDA TEACHERS' EDUCATION CONSULT (UTEC)

### Uganda Certificate of Education

#### **CHEMISTRY**

Paper 2

2 hours

#### INSTRUCTIONS TO CANDIDATES:

Section A consists of 10 structured questions. Answer ALL questions in this section. Answers to the questions must be written in the spaces provided.

Section B consists of 4 semi – structured questions. Answer only two questions from this section. Answers to the questions must be written in the answer booklets provided.

In both sections all working must be shown clearly.

Where necessary

$$[Zn = 65, O = 16, H = 1, Pb = 207]$$

1 mole of a gas occupies 22.4 dm<sup>3</sup> at s.t.p

1 mole of a gas occupies 24dm<sup>3</sup> at room temperature

	For Examiner's use only													
1	2	3	4	5	6	7	8	9	10	11	12	13	14	TOTAL

© 2022 UTEC Mock Examinations

Turn Over



# SECTION A: (50 MARKS)

## Answer ALL questions in this section

(a)	State	e what was observed.	(½ ,mark)
••••			
(b)	(i)	Name the piece of apparatus that can be used to separate the r	
			(½ mark)
••••	(ii)	Give a reason for your answer in (b) (i) above.	(½ mark)
•••••		•••••••••••••••••••••••••••••••••••••••	
(c)	sodiı	other experiment, ammonium chloride was heated as a mixture van chloride.	with
	State (i)	What was observed during the process of heating?	(01 1)
•••••	•••••		(01 mark)
•••••			•••••
	(ii) <sup>‡</sup>	The practical application of heating the mixture.	( ½ mark)
(d)		ion one example of a mixture which can be separated by fraction	
	distil	lation.	(01 mark)
	•••••		: 1
The a	atomic r	numbers of elements, $X$ , $Y$ and $Z$ are 6, 12 and 17. Respectively: the electronic configuration for:-	
	(i)	X:	(½ mark)
	(ii)	Y:	(½ mark)
	(iii)	Z:	(½ mark)

2.

	(b)	Writ (i)	e the formula for the product when:  X, burns in the excess oxygen.	(01 mark)
٠.,			39407 J	
		(ii)	Atoms of Y are burnt in the molecules of Z.	(01 mark)
			24.	47
	(c)		r was added to the compound in (b) (i) and the pro-	oduct tested with litmus
		(i)	What was observed?	(01 mark)
	•••••	•••••	······································	
	•••••	(ii)	Any one conclusion that can be drawn following yo	3 3
	•••••		······································	
3.	(a)		um peroxide reacts with water to produce oxygen according equation.	ording to the
		$2Na_2$	$O_2^{\frac{1}{2}}(s) + 2\dot{H}_2O(l) \longrightarrow 4 NaOH(aq)$	$+ O_2(g)$
		(i)	State what is observed during the reaction.	(01 mark)
	•••••	••••••		
		(ii)	Calculate the volume of oxygen gas produced at r 3.12g of sodium peroxide were completer reacted w	coom temperature when
	. :		$(Na = 23, 0 = 16, 1 \text{ mole of a gas occupies } 24\text{dm}^3$ at room temperature	m <sup>3</sup> at room temperature. e. (02 marks)
	••••••	••••••	,,	
		• • • • • • • • •		***************************************
	•••••	•••••		
	•••••	••••••		
	•••••	••••••		

(b)	A small sample of burning sulphur was plunged into a gas jar some oxygen gas. State:	
(i)	What was observed?	(01 mar)
(ii)	The effect of the product(s) in (c) (i) on litmus paper(s).	(01 mar
••••		
Dilut	e hydrochloric acid was added until in excess to zinc granules and t	he gas evolv
was p	bassed over heated lead (II) oxide as shown in the figure 1. Below.	
	25	
	Dilute hydrochloric acid	
	e e e e e e e e e e e e e e e e e e e	
	Combustion tube	
	psnq (II)	
/	J   Ç Ç Ç	
		5
	Heat U - tube	
		70
	Zinc granular Fused calcium	
(a)	State;	
	(i) What was observed in the combustion tube?	(01 mar
	***************************************	
<i></i>	(ii) The role of fused calcium chloride in the U-tube.	(½ mar

(b)	(i)	between zine and dilute hydrochloric acid.	(1 ½ mark)
••••			
	(ii)	that takes place in the combustion tube.	(1 ½ marks)
(c)		one other oxide besides Lead (II) oxide that could oustion tube.	d have been used in the (½ mark)
		npound, Q, reacts with dilute sulphuric acid to produce wing properties.	uce a colourless gas that
(i) (ii)		as acidified potassium dichromate solution from orange ches moist blue flowers to white.	e to green.
(a)	Nam		
	(i)	Compound Q.	( ½ mark)
		The colourless gas.	( ½ mark)
(b)	Writ	te an equation for the reaction between Q and dilute su	ia ,
	• • • • • • • • • • • • • • • • • • • •		
(c)	Stat		
	(i)	the practical application of the reaction when the a acidified potassium dichromate solution.	gas was bubbled through (½ mark)
· · ·			
···			

	(d)	The bl	eaching action of the colourless gas is by reduction. Write eaching action of the colourless gas is by reduction. Write each to show how the gas reduces moist blue flowers.	quation for the (01 mark)
				•
6.	Wher in a p	n magne lastic be	sium powder was added into 25cm <sup>3</sup> of 0.2M solution of copperator, some heat was evolved.	er (II) chloride
	(a)	State; (i)	What else was observed apart from the evolution of heat?	(01 mark)
		.,,		
1:-		(ii)	Why the reaction liberates heat.	(01 mark)
	(b)		an ionic equation for the reaction that took place.	(01 mark)
			•••••	
: <u>2: 1</u>	,			
,,,,,,	(c)	Calcu	late the;	
		(i)	Number of moles of copper (II) ions contained in the 25 solution.	5.0cm <sup>3</sup> of the (01 mark)
	hes			, ,
	••••	• • • • • • • • • • • • • • • • • • • •		*********
d : :				
		(ii)	The mass of copper that was displaced by one mole of magnet of Magnesium were used in the reaction. ( $Cu=64$ , $Mg=24$ ).	esium if 1.92g (02 marks)
		•••••••		
	• • • • • • • • • • • • • • • • • • • •			••••

The diagram in the Figure 2 below shows the set up of apparatus that was used to carry out electrolysis of different electrolytes in the laboratory. Carbon electrodes were used in the experiment.




Fig. 2

	Name the substance that was formed at electrode, A, if the following e were used.  (i) Aqueous silver nitrate solution.	(½ mark)
	(ii) Copper (II) chloride solution in dilute form.	(½ mark)
(b)	Write an equation for the electrode reaction in (a) (i).	(01 mark)
(c)	If the electrolyte used was concentrated sodium chloride solution, product(s) that was formed at electrode:	
	(i) A:	( ½ mark) (½ mark)
(d)		(01 mark)

	(e)	If the electrodes were replaced with Copper electrodes and the electrodes (ii) done for some time, state what was observed at A.	(½ mark)
8.	(a)	State what would be observed if each of the following reagents was add	
	(i)	solution of aluminium chloride.  Lead (II) nitrate solution.	( ½ mark)
			•
	(ii)	Ammonia solution dropwise until in excess.	(01 mark)
	(b)	Write an equation for the reaction that took place in (a) (i) above.	(01 mark)
			•
	(c)	To the resulting mixture in (a) (ii) was added sodium hydroxide solution until in excess.	on dropwise
	(i)	State what was observed.	(01 mark)
, : :	(ii)	Give a reason to support your answer in (c) (i).	 (01 mark)
9.	(a)	In the presence of a catalyst, glucose can be converted to ethanol.  (i) Name the catalyst used.	(½ mark)
		(ii) Write equation for the reaction in which glucose is converted	

(b)	The eth	nanol obtained can be converted to carbon dioxide gas accordin	g to the
(0)		ing equation:	B 10 1110
	Calcula	$H(l) + 3O_2(g) \longrightarrow 2CO_2(g) + 3H_2O(l) + Heat ate the mass of ethanol that was burnt in oxygen to produce 544 2, O = 16, H = 1; 1 mole of ethanol burns to produce 1360KJ or$	.0KJ of heat f heat) (02 marks)
••••			
	•••••	***************************************	
(c)	State v		
	(i)	is formed when ethanol is heated with concentrated sulphuric a	icia. (½ mark)
			(/2 11111111)
	(ii)	is observed when the product in (c) (i) is treated with bromine.	(01 mark)
When	n concer	ntrated hydrochloric acid was added to manganese (IV) oxide an	d the mixture
		was evolved.	
(a)	(i)	Name the gas that was evolved.	(½ mark
			/1 1/ monica
		, Write equation for the reaction that took place.	(1 ½ marks)
(b)	State	what is observed when the gas is bubbled through:	(01 mark
	(i)	Cold dilute potassium hydroxide solution.	•

10.

	(ii)	A beaker containing moist red flowers.	( ½ mark)
	The c	as in (a) (i) was bubbled through a solution of sodium iodic	le until no further
(c)	the g	•	
	(i)	Write equation for the reaction.	(1 ½ marks)
	•••••		
		a drawn from the ec	mation you have
	(ii)	Suggest any conclusion that can be drawn from the edwritten in (c) (i).	(1 mark)
			••••
		SECTION B (30 MARKS)	
		Answer any TWO questions from this section	
Colo	um ovid	le is regarded as a hygroscopic substance.	
(a)	(i) \	Define the term a hygroscopic substance.	(01 mark)
( <i>a</i> )	• • • • • • • • • • • • • • • • • • • •	Give one other example of a hygroscopic substance.	(½ mark)
		State one laboratory use of hygroscopic substance.	(01 mark)
(b)		A, is mainly dried in the laboratory using calcium oxide.	(12 22
	(i)	Identify gas A.	( ½ mark)
	(ii)	Write an equation for the reaction that leads to the formati	
•	()	in the laboratory.	(1 ½ marks)
(c)	Exces	s water was added on to calcium oxide until there was no fu	
	(i)	Write equation for the reaction that took place.	( 1 ½ marks)
	(ii)	Using equations, explain the changes that would result	,
		gas was bubbled through the product(s) in (c) (i) until in e	
			(4 ½ marks)
(d)	State;		
	(i)	The conditions for the reaction of calcium oxide with nitri	c acid.
	* ********	Write equation for the reaction that takes place.	(2 ½ marks)
	(ii)	Why the reaction in (d) (i) is not possible with sulphuric a require).	ecid. (Equation not (02 marks)

įσ.

11.

12.		rogen chloride gas can be prepared in the laboratory by react al and an acid.	
	(a)	(i) Name the acid commonly used for this preparation.	(½ mark)
	( )	(ii) State the conditions for the reaction.	(1 mark)
		(iii) Write equations for the reaction.	$(1 \frac{1}{2} \text{ marks})$
	(b)	Iron fillings when heated in dry hydrogen chloride react accor	ding to the following
	(-)	equation:	
		$Fe(s) + 2HCl(g) \longrightarrow FeCl_2(s) + H_2(g)$	
		(i) Calculate the volume of hydrogen chloride gas that	would be required to
,		react with iron at s.t.p to produce 5.0g of Iron (II) chlor	ride.
		(Fe = $56$ , Cl = $35.5$ ; 1 mole of a gas occupies $22.4$ dm	<sup>3</sup> at s.t.p).
			(2 ½ marks)
		(ii) The anhydrous Iron (II) chloride formed in (b) was	dissolved in water.
		State what was observed and write an ionic equation for	or the reaction.
			(2 ½ marks)
	(c)	Dilute sodium hydroxide solution was added to the solution	(product) in (b) (ii)
	(0)	drop wise until in excess.	
		(i) State what was observed.	(01 mark)
		(ii) Write an ionic equation for the reaction.	(1 ½ marks)
	(d)	Explain;	
	(u)	(i) The effect of bubbling hydrogen chloride gas through	h a solution of silver
		nitrate.	(2 ½ marks)
			`
		(ii) Why aqueous hydrogen chloride liberates hydrogen whereas hydrogen chloride in methyl benzene does no	
		whereas hydrogen chloride in memyr benzene does no	(2 /2 marks)
12	Midwa	gen can react with hydrogen in the Haber process to produce an	
13.			imona gas.
	(a)	State the;	(01 mark)
		(i) Source of nitrogen and hydrogen	•
		(ii) Factors that can affect the yield of ammonia in the Hal	
			(1 ½ marks)
	(b)	Write an equation for the reaction leading to the formation	
		Haber process.	(01 mark)
	(c)	Write an equation for the reaction to show oxidation of amme	
		(i) by Copper (II) oxide	(1 ½ marks)
		(ii) in the presence of hot platinum.	(1 ½ marks)
	(d)	Using equations, outline how the product(s) in (c) (ii) can	be converted to nitric
	(47)	acid.	(3 ½ marks)

Ammonia gas can react with concentrated sulphuric acid to produce ammonium (e) sulphate according to the following equation:  $2NH_3(g) + H_2SO_4(l) \longrightarrow (NH_4)_2SO_4(S)$ Calculate the mass of ammonium sulphate that would be produced by 0.96 dm3 of ammonia gas at room temperature. (N = 14, S = 32, O = 16, one mole of a gas occupies  $24 \text{dm}^3$  at room (02 marks) temperature. Explain why an aqueous solution of ammonium sulphate turns blue litmus (ii) (02 marks) paper red. When aqueous ammonia solution was added to a solution containing zinc ions, a (f) white precipitate Y which dissolved in excess ammonia to give a colourless solution was formed. ( ½ mark) Identify Y. (i) Write the formula for the cation in the colourless solution. ( ½ mark). (ii) (02 marks) State the difference between an acid and a salt. (a) Describe; (b) how a pure dry sample of lead (II) carbonate can be prepared in the (i) (04 marks) laboratory. (no diagram is required) (02 marks) the effect of heat on lead carbonate. Lead (II) carbonate reacts with dilute nitric acid according to the following (c) equation.  $PbCO_3(s) + 2HNO_3(aq) \longrightarrow Pb(NO_3)_2(aq) + CO_2(g) + H_2O(l)$ Calculate the mass of lead (II) carbonate that is required to react completed with  $200 \text{cm}^3$  of 0.2M dilute nitric acid. (Pb = 207, C = 12, O = 16). (2 ½ marks) State what would be observed if into a solution of lead (II) ions was added; (e) ( ½ mark) Drops of potassium iodide. (i) Ammonia solution drop wise until in excess. (01 mark) (ii) Dilute hydrochloric acid and the mixture heated then allowed to cool. (iii) (1 ½ marks) Write an equation to illustrate your answer in (d) (ii) above. (1 ½ marks)

14.

Û