Name	 Centre / Index No/
School	 Signature
P525/3	
CHEMISTRY (PRACTICAL)	

WAKISSHA

Uganda Advanced Certificate of Education

CHEMISTRY PRACTICAL

Paper 3

3 hours 15 minutes

Instructions to Candidates:

Answer all questions.

Paper 3 3¹/₄ hours

- Record your answers on this question paper in the spaces provided.
- Mathematical tables and silent non-programmable calculators may be used.
- Reference books (i.e, textbooks, books on qualitative analysis etc.) should **not** be used.
- Candidates are **not** allowed to start working with the apparatus for the first 15 minutes.

 This time is to enable candidates to read the question paper and make sure they have all apparatus and chemicals that they may need.

	For Exam	miners' Use Only	W W
Q.1	Q.2	Q.3	Total
1			
			1 - 3/2/2 point 37 a

You are provided with the following. GA1; which is approximately 0.1M sodium thiosulphate solution. GA2; which is a solution prepared by dissolving 1.0g of Potassium iodate in distilled water. Liquid E; which is a commercial bleaching agent containing sodium hypoch GA3; which is 0.5M potassium iodide GA4; which is 1.0M sulphuric acid	
You are required to: (i) Standardize GA1 using GA2 (ii) Determine the percentage of chlorine in the commercial bleaching age	ent.
Theory	
In acidic medium, Iodate ions react with Iodide ions according to the equation $IO_{3(aq)}^{-} + 6H_{(aq)}^{+} + 5I_{(aq)}^{-} \longrightarrow 3I_{2(aq)}^{-} + 5H_{2}O_{(l)}^{-}$	on
In acid medium hypochlorite ions produce chlorine according to the following $Cl^{-}_{(aq)} + OCl_{(aq)} + 2H^{+}_{(aq)} \longrightarrow H_2O_{(I)} + Cl_{2(aq)}$	ng equation.
The chlorine produced displaces Iodine from Potassium Iodide $Cl_{2(aq)} + 2I_{(aq)} \longrightarrow I_{2(aq)} + 2Cl_{(aq)}$	
Procedure A Pipette 10.0cm³ of GA2 into a conical flask and add 10.0cm³ of GA4 follow of GA 3. Titrate the liberated Iodine with GA1 from the burette until the mixture is particular that the description of the start and continue with the titration until the end point. Repeat the titration until you obtain consistent results. Record your result in the table 1 below.	
Volume of pipette used cm ³	$(0\frac{1}{2} \text{ mark})$
Table 1	
Final burette reading (cm ³)	<i>a</i>
Initial burette reading(cm ³)	
Volume of GA1 used(cm ³)	
Titre value of GA1 used to calculate the average volume.	(04½ marks) (½ mark)
Average volume GA1	(02½ mark)

deulate the molar concentration	on or GAT.		(6 mark)
		2	
	×		
		=	
roducer B Measure 30.0cm ³ of liquid E in the solution GA5.			
Measure 30.0cm ³ of liquid E ir	conical flask. Add GA1 from the bur nd continue with the	10cm ³ of GA4 followed bette until the mixture is particular titration until it turns contact.	by 10.0cm ³ of alle yellow. blourless.
Measure 30.0cm ³ of liquid E in nark. Label the solution GA5 . ipette 10.0cm ³ of GA5 into a GA3 . Sitrate the liberate Iodine with add 1cm ³ of starch indicator a Repeat the titration two more the Results	conical flask. Add GA1 from the bur nd continue with the	10cm ³ of GA4 followed bette until the mixture is particular terms of the titration until it turns coordinate results in table II below	by 10.0cm ³ of alle yellow.
Measure 30.0cm ³ of liquid E in mark. Label the solution GA5. ipette 10.0cm ³ of GA5 into a GA3. Pitrate the liberate Iodine with add 1cm ³ of starch indicator a Repeat the titration two more to Results	conical flask. Add GA1 from the bur nd continue with the	10cm ³ of GA4 followed bette until the mixture is particular terms of the titration until it turns coordinate results in table II below	by 10.0cm ³ of alle yellow. blourless.
Measure 30.0cm ³ of liquid E in mark. Label the solution GA5. ipette 10.0cm ³ of GA5 into a GA3. Citrate the liberate Iodine with Add 1cm ³ of starch indicator a Repeat the titration two more to Results Volume of pipette used	conical flask. Add GA1 from the bur nd continue with the	10cm ³ of GA4 followed bette until the mixture is particular terms of the titration until it turns coordinate results in table II below	by 10.0cm ³ of alle yellow. blourless.
Measure 30.0cm ³ of liquid E in mark. Label the solution GA5. ipette 10.0cm ³ of GA5 into a GA3. Titrate the liberate Iodine with Add 1cm ³ of starch indicator a Repeat the titration two more to Results Volume of pipette used	conical flask. Add GA1 from the bur nd continue with the	10cm ³ of GA4 followed bette until the mixture is particular terms of the titration until it turns coordinate results in table II below	by 10.0cm ³ of alle yellow. blourless.
Measure 30.0cm³ of liquid E in mark. Label the solution GA5. lipette 10.0cm³ of GA5 into a GA3. Citrate the liberate Iodine with Add 1cm³ of starch indicator a Repeat the titration two more to Results Volume of pipette used	GA1 from the burnd continue with the imes and record you	10cm³ of GA4 followed bette until the mixture is partial than the titration until it turns concur results in table II below cm³	by 10.0cm ³ of alle yellow. blourless.

Question

Number of moles Iodine liberated by 10.0cm ³ of GA5	(01mark)
Number of moles found instructed by Toloria of office	M
Number of moles of aqueous chlorine in 30cm ³ of E	(02½ marks)
•	
	11 11
	19 19 19 19 19 19 19 19 19 19 19 19 19 1
The mass of aqueous chlorine in 30cm^3 of E. ($Cl = 35.5$)	(02 marks)
Percentage of mass of aqueous chlorine in E.	(01 mark
Percentage of mass of aqueous emornic in E.	
200 m 1 A 2 Mars	
)	

You are provided with substance W, Which contains three cations and one anion. You are required to carry out the following tests on W, to Identify the cations and anion in W. Identify any gase(s) evolved.

Record your observations and deductions in the table below.

Tests	Observations	Deductions
 Heat a spatula endful of W strongly in a dry test tube. 		
endfuls of W with about 4cm³ of water, add dilute sodium hydroxide solution to the mixture drop- wise		
until in excess. Warm and filter. Keep both the filtrate and the residue.		
c) To the filtrate, add dilute nitric acid drop wise until the solution is just acidic. Divide the acidic solution into six parts		
(i) To the first part of the solution, add dilute sodium hydroxide solution drop wise until in excess. Warm the mixture.		
(ii) To the second part of the solution, add 2-3 drops of Potassium Iodide solution.		
(iii) To the third part of the solution add dilute ammonia solution drop wise until in excess.		
(iv) Use the fourth part of the solution to carry out a test of your own choice to identify one of the Cations in W.		
	The second secon	
		Turn-O

(v) To the fifth part of the solution, add 2-3 drops of lead(II) nitrate solution.		
(vi) Use the sixth part to carry out a test of your own choice to confirm the anion in W		
d) Wash the residue with water and dissolve it in dilute hydrochloric acid. Divide the solution into three parts.		
(i) To the first part of the solution, add dilute sodium hydroxide solution drop wise until in excess.		
(ii) To the second part of the solution, add ammonia solution drop wise until in excess		
(iii) Use the third part carry out a test of your own choice to confirm one of the Cations in W.		
e) (i) The anions in W	is	
(ii) Cations in W		and

You are provided with substance Q which is an organic compound. You are required to determine the nature of Q. Carry out the following tests on Q. and record your observations

and deductions in the table below.

3.

Test	Observation	Deductions
a) Burn a spatula endful of Q on a porcelain dish or spatula end.		
b) To about 3cm³ of Q in a test tube, add, about 3cm³ of water, shake and allow to stand. Then test the mixture with litmus paper. Divide the resultant mixture in to 5 portions.		
(i) To the first portion of the mixture add 3-4 drops of neutral iron (III) chloride solution.		
(ii) To the second portion of the mixture, add half a spatula endful of solid sodium hydrogen carbonate.		
(iii) To the third portion of the mixture add 2 - 3 drops of 2, 4 - dinihophenyl hydrazine.		
(iv) To the forth portion of the mixture, add an equal volume of ethanol followed by 2-3 drops of concentrated sulphuric acid. Heat the mixture and allow to stand. Then pour into a small beaker of water		
(v) To the fifth portion of the mixture, add 3-4 drops of acidified potassium dichromate solution and heat.		

c) To about 1cm ³ of silver		
nitrate solution in		
boiling tube, add an	Mark the second Ch	
equal volume of		
sodium hydroxide		
solution followed by	di e	
dilute ammonia		
solution until the		an end of the
precipitate formed just	Part	A company of the comp
dissolves. Then add		100 100
about 1cm3 of Q and	* × × ×	C1 41C 201 M
boil in water bath for		4 m 2 A-11
about 5-minutes. Allow		
to cool under running		
tap water.		

d) Comment on the nature of Q

END