P 510/1 **PHYSICS** Paper 1 UGANDA ADVANCED CERTIFICATE OF EDUCATION **PHYSICS** S.6 PAPER 1 - 2014 2 hours 30 minutes. **INSTRUCTIONS TO CANDIDATES:** Attempt **five** questions in all. Non-programmable scientific electronic calculators may be used.

= 9.81 ms⁻²

Assume where necessary.

Acceleration due to gravity, g

Stefan's Boltzmann's constant, σ	=	$5.7x10^{-8} \ Wm^{-2} \ K^{-4}$	
Radius of the earth	=	6.4 x 10 ⁶ m	
Radius of the sun	=	7.0 x 108m	
Radius of Earth's orbit about the sun	=	1.5 x10 ¹¹ m	
Speed of light in a vacuum, c	=	3.0 x 10 ⁸ ms ⁻¹	
Specific heat capacity of water	=	4,200J kg ⁻¹ K ⁻¹	
Universal gravitational constant, G	=	6.67 x10 ⁻¹¹ N m ² kg ⁻²	
Density of water	=	1000 kgm ⁻³	
Gas constant, R	=	8.31 J mol ⁻¹ K ⁻¹	
SE	CTION A		
3. (a) (i) State the law of conservation of e	energy.	(1)
(ii) A particle is projected at an acute show that the mechanical energy of the part	_	_	ir resistance
motion		(5)	
(b) A particle, hanging from a spring at reparticle is given a small vertical displacement motion and find the period.			
(c) A mass of 500 g is released from rest onto a scale pan of negligible mass hung from			
(i) the position of the scale pan when i	t first comes	to instantaneous rest	(5)

= 5.97 x 10^{24} kg

Mass of the earth

(ii) the amplitude of the subsequent motion of the particle	(2)
(iii) the period of the motion.	(2)
2. (a) Define the terms: <i>elastic limit</i> and <i>Young's modulus.</i>	(2)
(b) Explain the energy transformations that occur when a wire is stretch	ned
(i) elastically	(3)
(ii) plastically	(2)
(c) A rod of mild steel of uniform cross sectional area $0.0030~\text{m}^2$ and less stretched steadily until it breaks.	ngth 1.0 m is
(i) Sketch a graph to show the relationship between the force and the	he extension.
Explain the shape of the graph.	(4)
(ii) When the applied force on the rod is 120 kN the strain is 4.0×10^{-1}) ⁻⁴ .
Calculate Young's modulus for steel.	(3)
(iii) The rod is found to break at a stretching force of 240 kN.	
Explain why the stress at the section of the rod where the break much greater than 80000 kN $\mbox{m}^{\mbox{-}2}$.	occurs is likely to be
(iv) Just before the rod breaks, its extension is about 4 cm.	
Estimate the work done in stretching the rod by this amount. Sug- calculated value of the work done is likely to be less than the actual value.	ggest why the (4)
3. (a) (i) State Newton's law of gravitation.	(1)
(ii) Deduce the dimensions of the Universal gravitational constant	(2)

(b) Assume that a planet moves with uniform speed in a circular consumation sum. The periodic time of the planet is T. Show that the quantity (r^3/T^2) mass of the planet.	
(c) An artificial satellite is launched in a circular orbit round the ec \times 10 7 m above the surface of the Earth.	quator at a height of 3.60
(i) Determine from first principles the speed with which the sat to maintain it in the orbit.	cellite must be launched (5)
(ii) Determine the periodic time of the satellite	(3)
(iii) What deduction can be made from the result obtained in (ii) the satellite?) above with reference to (2)
(iv) If another satellite moves in a circular orbit concentric with satellite, at a height of 1.48 x 10^7 m above the surface of the Earth, wh this second satellite.	
4. (a) Distinguish between <i>scalar</i> and <i>vector</i> quantities. Give two ex	camples of each.(3)
(b) (i) Define the terms <i>time of flight</i> and <i>range</i> as applied to proj	ectile motion. (2)
(ii) A projectile is fired in air with a speed u at an angle θ to the of flight of the projectile.	horizontal. Find the time (2)
(c) State the conditions for the equilibrium of a rigid body under t forces.	he action of coplanar (2)
(d) A mass of 5.0 kg is suspended from the end A of a uniform beal length 1.0 m. The B of the beam is hinged in the wall. The beam is kep attached to the mid-point of the beam and to a point C, in the wall at Calculate	t horizontal by a rope
(i) the tension in the rope	(6)

(ii) the force exerted by the hinge on the beam.	(5)
8. (a) (i) Distinguish between thermionic emission and photoelectric emission.(ii) Apart from them carrying negative charge, state four other properties of (2)	(3) cathode rays.
(b) Describe an experiment to show that cathode rays indeed carry negative ch	arge. (4)
(c) A beam of electrons moving with a velocity v enters normally a strong magn which is uniform and of induction B. If each electron has a mass m and charge e, s assumptions made,	
(i) derive an expression for the distance d covered by an electron while movifield.	ving in the (3)
(ii) write down the relationship between d and B	(1)
(d) A charged oil drop falls under gravity in air with a velocity of $4.90 \times 10^{-4} \text{ms}^{-1}$ parallel plates 5 mm apart. When a p.d of 4500 V is connected between the plate rises steadily with a velocity of $1.10 \times 10^{-4} \text{ms}^{-1}$.	
Assuming that the effect of buoyancy on the drop is negligible, calculate	
(i) the radius of the oil drop.	(3)
(ii) the number of electrons on the drop	(4)
(Density of oil = 900 kg m $^{-3}$; coefficient of viscosity of air =1.81 x 10^{-5} Nm $^{-2}$ s)	
1.(a) State the laws of solid friction	(3)
(b) (i) Describe an experiment to determine the co-efficient of friction between (4)	two surfaces

(ii) A particle of mass m is projected up a rough plane of ang maximum and slides back. Prove that when it again attains its or distance	·
<u>u²μsecø</u>	
$g(tan^2\phi - \mu^2)$	
down the plane from the point of projection, given that $\mu,$ the cotanø.	-efficient of friction is less than (6)
(c) (i) Define co-efficient of viscosity and laminar flow	(2)
(ii) Given that the volume of a liquid issuing per second from efficient of viscosity η , the radius of the pipe a and the pressure ϱ	
Volume per second = \underline{kpa}^4	
ηΙ	(5)
5. (a) (i) State Boyle's law.	(1)
(ii) Describe an experiment to verify Boyle's law.	(6)
(b) (i) Explain why the pressure of a fixed mass of gas in a closed rigid temperature of the container falls.	d container decreases when the
(ii) State the conditions for an isothermal change	(2)
(c) (i) Derive the equation of state. (ii) A ball of volume 4 litres at a temperature of 20°C contains nitro	(4) ogen at a pressure of

the tem	⁵ Pa. After some more nitrogen had been pumped in, the volume of perature 30° C and the pressure 5.5×10^{5} Pa. What mass of nitrogen nitrogen = 28 g		•
	Define surface tension and show that it is equal to the work done urface under isothermal conditions.	per unit area in i (3)	ncreasing the
	Find the work required to break up a drop of water of radius 0.5 cmm. [Surface tension of water is $2.27 \times 10^{-2} \text{ Nm}^{-1}$]	n into drops of w (4)	ater each of
(b) De	escribe an experiment to measure surface tension of a liquid by the	capillary tube m	ethod.
surface	Derive an expression for the excess pressure in a soap bubble of ratension γ. (The pressure inside an air bubble formed 40 cm below the water so	3)	solution has
	10 ⁵ Pa. If the surface tension of water is 2.27 x 10^{-2} Nm ⁻¹ find the rapheric pressure = 1.013 x 10^{5} Pa]	dius of the bubb (4)	le.
4. (a)	(i) Define the two principal molar heat capacities of a ga	zs.	(02)
	(ii) State, which one is greater, and explain why this is s	60.	(02)
molar	(iii) Show that the difference between the molar heat ca gas constant.	<i>pacities</i> is equ	ual to the
(b)	State the conditions for an adiabatic process.		(02)

adiabatically to twice its original volume and then is compressed to its original volume. [Take ratio of the principal molar heat capacities, γ	isothermally		
(i) Sketch and label the two stages on a P-V diagram.	(02)		
(ii) Find the <i>final pressure</i> and <i>temperature</i> of the gas.	(07)		
7. (a) State Prevost's theory of exchange with reference to heat radiation. (1)			
(b) Explain why metals are better conductors of heat compared to wood. (4)			
(c) A is a composite slab made up of two layers of thickness d_1 and d_2 with thermal conductivities k_1 and k_2 respectively. B is a slab of thickness $d_1 + d_2$ with thermal conductivity k. One face of each of the slabs is maintained by heating at a temperature θ_1 and the other end at θ_2 . If the rate of conduction per unit area through the two slabs is the same, find k in terms k_1 , k_2 , d_1 and d_2 .			
(d) The heat radiation received by the earth from the sun is 1.4×10^3 Wm $^{-2}$. Assuming this 90% of what the sun emits as a black body, estimate the temperature of the sun. (4)			
(e) With the aid of a labeled diagram, describe the construction and mode of operation pyrometer. (5)	of the optical		
5 (a) (i) Define thermal conductivity.	(1)		
(ii) Explain the mechanism of heat transfer in metals.	(3)		

(b) (i) What is a black body?	(1)
(ii) Draw sketch graphs to show the variation of relative intensity of black b	ody
radiation with wavelength for different temperatures.	(2)
(iii) Explain the appearance of a metal ball placed in a dark room when its terprogressively raised from room temperature to just below melting. (3)	mperature is
(c) A heating element in form of a cylinder 30 cm long and 1.5 cm in diameter output of 1.8 kW . If its radiation is 85% that of a black body, find its tempera (4)	
6. (a) What assumptions are necessary in the derivation of the Kinetic theory the pressure of an Ideal gas? (4)	expression for
(b) A beam of 2 x 10 ²² Nitrogen atoms, each of mass 2.32 x 10 ⁻²⁶ kg is incident a wall of a cubical container of edge 5.0cm . The beam is reflected through 18 0 mean speed of the atoms is 500 ms ⁻¹ , find the pressure exerted by the Nitrogen (4)	0 °. If the
(c) (i) State Dalton's law of partial pressures.	(1)

(iii) Describe an experiment to determine the thermal conductivity of a metal. (6)

	(ii) Two containers A and B of volume 3 x 10 ³ cm³ and 6 x 10 ³ cm³ respectively. Container B is cooled to 273 K . Find the final pressure of the B (5)	r A is heate	ed to
	(i) Use the kinetic theory of gases to explain the effect of increasing temperation vapour pressure. (3)	erature oi	ı
	(ii)Sketch a pressure versus volume curve for a real gas undergoing compitits critical temperature. Explain the main features of the curve.	ression belo 3)	ЭW
7.	(a) (i) Define specific latent heat of fusion.	(1)	
	(ii) Describe an experiment to determine the specific latent heat of fus (5)	sion of w	ater.
	(b) (i) Explain why the molar heat capacity of an ideal gas at constant promethat at constant volume.	pressure di 3)	ffers
	(ii) Derive an expression for the difference of the above molar heat ca	pacities. 4)	
	(c) (i) State the conditions necessary for a reversible adiabatic process.	(2)	

	(ii) A gas initially occupying a volume of 1.0 litre at a temperature of 47 % pressure of 1.40 x 10 5 Pa expands isothermally to a volume of 2.0 litres . It compressed adiabatically to its original volume. If $\gamma = 1.4$ find the final team of pressure. (5)	t is then
7.	(a) (i) The specific heat capacity of copper is 400J kg ⁻¹ K ⁻¹ . Explain the mean statement.	ing of the (1)
	(ii) State the factors that determine the rate of cooling of a body.	(2)
	(iii) Explain why a small body cools faster than a bigger one of the same mashape.	aterial and (4)
minim (7)	(b) With the aid of a well labeled diagram, describe an electrical method o the specific latent heat of vaporization of water, stating the precautions ta nize heat losses	_
	(c) Two taps, one delivering water at a temperature of 25°C at a rate of 0 .3 other delivering water at a temperature of 15°C at a rate of 0.4 g s ⁻¹ , are di well-lagged copper calorimeter of mass 50 g containing ice at 0°C while stimulates the contents of the calorimeter are found to have attained a temperature.	rected to a rring. After 5

5°C.

SECTION C

Find the mass of ice originally in the calorimeter. Specific latent heat of fusion of ice = $3.36 \times 10^5 \, \text{Jkg}^{-1}$

8. (a) (i) Sketch the I-V characteristic for gaseous conduction and explain the main features of the curve. (4)

(6)

- (ii) Explain why this is not a good method for producing cathode rays. (2)
 - (b) (i) What is meant by **thermionic emission**? (1)
 - (ii) Describe an experiment to show that cathode rays travel in straight lines. (4)
 - (c) A horizontal beam of electrons, moving with a uniform speed, enters a uniform electric field between two horizontal parallel charged plates. Show that the path of the beam between the plates is a parabola. (4)
 - (d) A charged oil drop of mass $6 \times 10^{-15} \, kg$ falls vertically in air with a steady velocity between two parallel vertical plates $5 \, mm$ apart. When a p.d of $3000 \, V$ is applied between the plates, the drop now falls with a steady velocity at an angle 58.5° to the vertical. Calculate the charge Q on the drop. (Neglect the up thrust on it)

P 510/1 **PHYSICS** Paper 1 UGANDA ADVANCED CERTIFICATE OF EDUCATION **END OF TERM 1 – 2014 PHYSICS S.6 PAPER 1** 2 hours 30 minutes. **INSTRUCTIONS TO CANDIDATES:** Attempt **five** questions in all, choosing at least one from each of the sections A, B and C. Assume where necessary.

= 9.81 ms⁻²

Acceleration due to gravity, g

Mass	of the earth	=	$5.97 \times 10^{24} \text{ kg}$	
Stefai	n's Boltzmann's constant, σ	=	5.7x10 ⁻⁸ Wm ⁻² K ⁻⁴	
Radiu	us of the earth	=	$6.4 \times 10^6 \text{m}$	
Radiu	us of the sun	=	$7.0 \times 10^8 \text{m}$	
Radiu	us of Earth's orbit about the sun	=	1.5 x10 ¹¹ m	
Speed	d of light in a vacuum, c	=	$3.0 \times 10^8 \text{ms}^{-1}$	
Speci	fic heat capacity of water	=	4,200J kg ⁻¹ K ⁻¹	
Unive	ersal gravitational constant, G	=	6.67 x10 ⁻¹¹ N m ² kg ⁻²	
Dens	ity of water	=	1000 kgm ⁻³	
Gas c	onstant, R	=	8.31 J mol ⁻¹ K ⁻¹	
	SEC	CTION A		
1(a)	Define the following:			
	(i) Tensile stress			(1)
	(ii) Tensile strain			(1)
	(iii) Young's modulus			(1)
(b)	(i) Sketch the stress-strain curve for	a copper w	ire and explain the characteristic	
	features of the curve.			(4)
	(ii) How does the curve in b(i) compa	re with tha	t of a material like glass?	(2)


(c)	Derive an expression for the energy stored in a unit volume of a stretched elastic material, expressing your answer in terms of Young's modulus and the strain of the		
	material.	(4)	
(d)	A cylindrical copper rod of length 0.5 m and diameter 2 cm is fixed between two rigid supports at a temperature of 20°C. The temperature of the rod is raised to 70°C Calculate	Ξ.	
	(i) the force exerted on the rigid supports, at this temperature.	(4)	
	(ii) the energy stored in the rod at 70°C	(3)	
	Young's modulus for copper = $1.2 \times 10^{11} \text{ Nm}^{-2}$ (Mean coefficient of linear expansion copper between 20°C and 70°C is $1.7 \times 10^{-5} \text{ K}^{-1}$)	ı of	
2(a)	Define simple harmonic motion.	(1)	
(b)	A particle moves so that its displacement x after time t is given by $x = r \sin 2\pi t$		
	(i) Show that the particle executes simple harmonic motion	(3)	
	(ii) Using the same axes sketch graphs of displacement against time, velocity against time and acceleration against time.	(3)	

(c)	submerged. Show that if the cylinder is given a small vertical displacement and ther released, it executes simple harmonic motion of period $\pi \sqrt{\left(\frac{3h}{g}\right)}$	
		(6)
(d)	A cube of mass 100 g rests on a light scale pan that is supported on a vertical sprin constant $24.5~\rm Nm^{-1}$. The pan is then set to oscillate vertically. Find the maximum	ıg of
	amplitude for the cube to remain in contact with the pan throughout the motion.	(3)
(e)	(i) Distinguish between free and damped oscillations and sketch a displacement-t graph for each	ime (4)
3(a)	(i) State stokes' law of viscosity	(1)
	(ii) A ball of density σ and radius r is released to fall through a liquid of density ρ viscosity, η . Derive an expression for its terminal velocity during its motion.	and (3)
	(iii) Two identical rain drops are observed to fall with a terminal velocity of 0.2 r If the two coalesce into one drop, what will be the new terminal velocity?	ms ⁻¹ (4)
(b)	(i) State Bernoulli's principle	(2)

(ii) Explain one application of Bernoulli's principle

(3)

(c) The diagram below shows a pipe delivering water



The cross-sectional area at section A is 30 cm² and the velocity there is 4 ms⁻¹ while the cross-sectional area at B is 15 cm². Find:

(ii) the static pressure at B if the atmospheric pressure is
$$1 \times 10^5 \text{ Nm}^{-2}$$
. (5)

4(a) (i) Show that in uniform circular motion the accelerating force is towards the centre of the circle and derive an expression for the acceleration. (4)

(2)

- mass M in an orbit of radius R. (3)

(b)

(iii) Explain the effect of friction between such a satellite and the atmosphere in which it moves. (4)

SECTION B

5(a) Describe an experiment to verify Newton's law of cooling. (5)

(b) (i) Distinguish between *a real* and *an ideal* gas. (3)(ii) Derive the expression: $P = \frac{1}{3}\rho \overline{c^2}$ for the pressure of an ideal gas of density ρ and mean square speed $\overline{c^2}$. (6) (c) (i) Explain why the pressure of a fixed mass of gas in a closed container increases when temperature of the container is raised. (2)(ii) Nitrogen gas is trapped in a container by a movable piston. If the temperature of the gas is raised from 0°C to 50°C at constant pressure of 4.0 x 10⁵ Pa and the total heat added is 3.0×10^4 J, calculate the work done by the gas. [The molar heat capacity of nitrogen at constant pressure is 29.1 J mol⁻¹ K⁻¹; C_p/C_v = 1.4] (4) 6(a) (i) What is meant by the term temperature? (1) (ii) Give two characteristics of a material to be chosen as a thermometric substance (2)(iii) Name six thermometric properties that can be used to measure temperature. (3)(b) Define a Celsius scale based on a fixed mass of gas at constant volume. (2) (c) (i) Explain briefly why thermometers based on different thermometric properties

give different values for the same temperature.


(1)

	(ii) A thermometer is constructed with a liquid that expands according to the relati	on
	$V_t = V_0(1 + \alpha t + \beta t^2)$	
	where V_t = volume of liquid at temperature $t^0 C$, V_0 = volume of liquid at $0^0 C$	
	α and β are constants such that $\alpha = 1000\beta$.	
	What will the liquid thermometer read when the gas thermometer reads 200°C?	(5)
(d)	(i) What is a thermo-junction?	(2)
	(ii) Sketch a graph of the emf of a thermocouple for the range before and after the neutral temperature.	(2)
	(iii) State two advantages and two disadvantages of thermocouple thermometers over other types.	er (2)
7 (a)	With the aid of diagram, discuss briefly variations of temperature gradient for a steam of the bar is;	eel
	(i) unlagged	(2)
	(ii) lagged	(2)
(b)	(i) In the determination of thermal conductivity of a poor conductor such as cork, substance is made thin and fairly of large cross-sectional area. Explain why this is s	
	(ii) Describe how the thermal conductivity of cork can be determined.	(5)

(c)	An aluminium pan of diameter 20.0cm and thickness 2.0mm is filled with water and heated such that the water boils producing steam at a rate of 5.0gs ⁻¹ at steady state conditions.	
	(i) Calculate the temperature of the external surface of the pan	(4)
	(ii) State the assumption made	(1)
(d)	(i) State Stefan's law of black body radiation.	(1)
	(ii)As a piece of charcoal is steadily heated up, it appears reddish in colour before turning whitish. Explain this observation.	(3)
	SECTION C	
8(a)	(i) Sketch the <i>I-V characteristic</i> for gaseous conduction.	(2)
	(ii) Explain the main features of the curve.	(3)
(b)	A beam of electrons moving with a velocity \mathbf{v} , enters normally a strong magnetic which is uniform and of induction \mathbf{B} . If each electron has a mass \mathbf{m} and charge \mathbf{e} , assuming that the electrons suffer a deviation of 180° by the time they leave the magnetic field,	ield,
	(i) derive an expression for the distance d , covered by an electron while moving in field.	the (3)

	(ii) write down the relationship between ${\bf d}$ and ${\bf B}$.	(1)
(c)	(i) Draw a labeled diagram of a cathode ray oscilloscope (C.R.O)	(3)
	(ii) Describe how an alternating p.d may be measured using a C.R.O	(3)
(d)	A horizontal beam of electrons, moving with a uniform speed, enters a uniform electric field between two horizontal parallel charged plates. Show that the path of the beam between the plates is a parabola.	(5)
9(a)	A high p.d is applied across two electrodes in air contained in a closed glass tube. Describe, with the aid of labeled diagrams, what will be observed when the pressi in the tube is progressively reduced down to very low pressures.	re (5)
(b)	(i) List four main properties of cathode rays.	(2)
	(ii) Describe an experiment to show that cathode rays carry a negative charge.	(4)
(c)	(i) In the Millikan's oil-drop experiment, why is there a need to surround the chamber with a constant temperature bath?	(2)
	(ii) Explain how Millikan's experiment proves that charge is quantized.	(2)

m was found to fall under gravity at a terminal velocity of 0.004 cm s⁻¹ and 0.012 cm s⁻¹ when a field of 2 x 10 ⁵ Vm⁻¹ was suitably applied. Given that the of the medium was 2.122 x 10 ⁻⁵ Nsm⁻² , determine the number of electrons		
on the drop.	Qn	Mks
(5)		
END		



P 510/1 PHYSICS

Paper 1

BEGINNING OF TERM TWO EXAMS – 2017

S.6 PHYSICS

Paper 1

2 hours 30 minutes.

INSTRUCTIONS TO CANDIDATES:

Attempt **five** questions in all, choosing at least one from each of the sections A, B and C. Indicate the questions attempted in the table in the RHS corner.

Assume where necessary.

Acceleration due to gravity, g = 9.81 ms^{-2}

Mass of the earth = $5.97 \times 10^{24} \text{ kg}$

Stefan	's Boltzmann's constant, σ	=	$5.7 \times 10^{-8} \ Wm^{-2} \ K^{-4}$	
Radius of the earth		=	6.4 x 10 ⁶ m	
Radiu	s of the sun	=	$7.0 \times 10^8 \text{m}$	
Radiu	s of Earth's orbit about the sun	=	$1.5 \text{ x} 10^{11} \text{m}$	
Speed	of light in a vacuum, c	=	$3.0 \times 10^8 \text{ms}^{-1}$	
Specif	ic heat capacity of water	=	4,200J kg ⁻¹ K ⁻¹	
Unive	ersal gravitational constant, G	=	$6.67 \text{ x} 10^{-11} \text{N m}^2 \text{ kg}^{-2}$	
Densi	ty of water	=	1000 kgm ⁻³	
Gas co	onstant, R	=	8.31 J mol ⁻¹ K ⁻¹	
		SECTION A		
1(a)	Define the following:			
	(i) Tensile stress			(1)
	(ii) Tensile strain			(1)
	(iii) Young's modulus			(1)
(b)	(i) Sketch the stress-strain curve f	or a copper wi	ire and explain the characteristic	
	features of the curve.			(4
	(ii) How does the curve in b(i) com	pare with tha	t of a material like glass?	(2)


(c)	Derive an expression for the energy stored in a unit volume of a stretched elastic material, expressing your answer in terms of Young's modulus and the strain of the	
	material.	(4)
(d)	A cylindrical copper rod of length 0.5 m and diameter 2 cm is fixed between two rigid supports at a temperature of 20° C. The temperature of the rod is raised to 70° C Calculate	C.
	(i) the force exerted on the rigid supports, at this temperature.	(4)
	(ii) the energy stored in the rod at 70°C	(3)
	Young's modulus for copper = $1.2 \times 10^{11} \text{ Nm}^{-2}$ (Mean coefficient of linear expansion copper between 20°C and 70°C is $1.7 \times 10^{-5} \text{ K}^{-1}$)	ı of
2(a)	Define simple harmonic motion.	(1)
(b)	A particle moves so that its displacement x after time t is given by $x = r \sin 2\pi t$	
	(i) Show that the particle executes simple harmonic motion	(3)
	(ii) Using the same axes sketch graphs of displacement against time, velocity against time and acceleration against time.	st (3)

A uniform cylinder of height h floats vertically in water with ¾ of its height (c) submerged. Show that if the cylinder is given a small vertical displacement and then released, it executes simple harmonic motion of period $\pi \sqrt{\left(\frac{3h}{a}\right)}$ (6) A cube of mass 100 g rests on a light scale pan that is supported on a vertical spring of (d) constant 24.5 Nm⁻¹. The pan is then set to oscillate vertically. Find the maximum amplitude for the cube to remain in contact with the pan throughout the motion. (3) (e) (i) Distinguish between free and damped oscillations and sketch a displacement-time graph for each.. (4) 3(a) (i) State stokes' law of viscosity (1)(ii) A ball of density σ and radius r is released to fall through a liquid of density ρ and viscosity, η. Derive an expression for its terminal velocity during its motion. (3)(iii) Two identical rain drops are observed to fall with a terminal velocity of 0.2 ms⁻¹ If the two coalesce into one drop, what will be the new terminal velocity? (4) (b) (i) State Bernoulli's principle (2)

(c) The diagram below shows a pipe delivering water

(ii) Explain one application of Bernoulli's principle

(3)



The cross-sectional area at section A is 30 cm² and the velocity there is 4 ms⁻¹ while the cross-sectional area at B is 15 cm². Find:

- (i) the volume of water flowing per second (2)
- (ii) the static pressure at B if the atmospheric pressure is $1 \times 10^5 \text{ Nm}^{-2}$. (5)
- 4(a) (i) Show that in uniform circular motion the accelerating force is towards the centre of the circle and derive an expression for the acceleration. (4)
 - (ii) State two differences between uniform circular motion and simple harmonic motion. (2)

	(iii) A cyclist is to turn a corner of a horizontal circular track of radius r where the coefficient of friction between the tyres and the track surface if μ . For no skiddin velocity, v , must fulfill the condition that	
	$v < \sqrt{\mu r g}$	(3)
(b)	(i) Define gravitational potential.	(1)
	(ii) Find an expression for the total energy of a satellite of mass m circling the earmass M in an orbit of radius R.	th of (3)
	(iii) Explain the effect of friction between such a satellite and the atmosphere in which it moves.	(4)
	SECTION B	
5(a)	Describe an experiment to verify Newton's law of cooling.	(5)
(b)	(i) Distinguish between <i>a real</i> and <i>an ideal</i> gas.	(3)
	(ii) Derive the expression:	

(ii) Explain briefly the action of a centrifuge.

(3)

- $P = \frac{1}{3}\rho \overline{c^2}$ for the pressure of an ideal gas of density ρ and mean square speed $\overline{c^2}$. (6)
- (c) (i) Explain why the pressure of a fixed mass of gas in a closed container increases when temperature of the container is raised. (2)
 - (ii) Nitrogen gas is trapped in a container by a movable piston. If the temperature of the gas is raised from 0° C to 50° C at constant pressure of 4.0×10^{5} Pa and the total heat added is 3.0×10^{4} J, calculate the work done by the gas.

[The molar heat capacity of nitrogen at constant pressure is 29.1 J mol
$$^{-1}$$
 K $^{-1}$; $C_p/C_v=1.4$] (4)

- 6(a) (i) What is meant by the term temperature? (1)
 - (ii) Give two characteristics of a material to be chosen as a thermometric substance (2)
 - (iii) Name six thermometric properties that can be used to measure temperature. (3)
- (b) Define a Celsius scale based on a fixed mass of gas at constant volume. (2)
- (c) (i) Explain briefly why thermometers based on different thermometric properties give different values for the same temperature. (1)
 - (ii) A thermometer is constructed with a liquid that expands according to the relation $V_t = V_0(1 + \alpha t + \beta t^2)$

	where V_t = volume of liquid at temperature t^0C , V_0 = volume of liquid at 0^0C	
	α and β are constants such that α = 1000 β .	
	What will the liquid thermometer read when the gas thermometer reads 200°C?	(5)
(d)	(i) What is a thermo-junction?	(2)
	(ii) Sketch a graph of the emf of a thermocouple for the range before and after the	
	neutral temperature.	(2)
	(iii) State two advantages and two disadvantages of thermocouple thermometers or other types.	er (2)
7 (a)	With the aid of diagram, discuss briefly variations of temperature gradient for a stabar when the bar is;	eel
	(i) unlagged	(2)
	(ii) lagged	(2)
(b)	(i) In the determination of thermal conductivity of a poor conductor such as cork, substance is made thin and fairly of large cross-sectional area. Explain why this is s	
		(2)
	(ii) Describe how the thermal conductivity of cork can be determined.	(5)

(c)	An aluminium pan of diameter 20.0cm and thickness 2.0mm is filled with water and heated such that the water boils producing steam at a rate of 5.0gs ⁻¹ at steady state conditions.	
	(i) Calculate the temperature of the external surface of the pan	(4)
	(ii) State the assumption made	(1)
(d)	(i) State Stefan's law of black body radiation.	(1)
	(ii)As a piece of charcoal is steadily heated up, it appears reddish in colour before turning whitish. Explain this observation.	(3)
	SECTION C	
8(a)	(i) Sketch the <i>I-V characteristic</i> for gaseous conduction.	(2)
	(ii) Explain the main features of the curve.	(3)
(b)	A beam of electrons moving with a velocity \mathbf{v} , enters normally a strong magnetic which is uniform and of induction \mathbf{B} . If each electron has a mass \mathbf{m} and charge \mathbf{e} , assuming that the electrons suffer a deviation of 180° by the time they leave the magnetic field,	ield,
	(i) derive an expression for the distance d , covered by an electron while moving in field.	the (3)

	(ii) write down the relationship between d and B .	(1)
(c)	(i) Draw a labeled diagram of a cathode ray oscilloscope (C.R.O)	(3)
	(ii) Describe how an alternating p.d may be measured using a C.R.O	(3)
(d)	A horizontal beam of electrons, moving with a uniform speed, enters a uniform electric field between two horizontal parallel charged plates. Show that the path of the beam between the plates is a parabola.	(5)
		()
9(a)	A high p.d is applied across two electrodes in air contained in a closed glass tube. Describe, with the aid of labeled diagrams, what will be observed when the press in the tube is progressively reduced down to very low pressures.	
(b)	(i) List four main properties of cathode rays.	(2)
	(ii) Describe an experiment to show that cathode rays carry a negative charge.	(4)
(c)	(i) In the Millikan's oil-drop experiment, why is there a need to surround the chamber with a constant temperature bath?	(2)
	(ii) Explain how Millikan's experiment proves that charge is quantized.	(2)

(d) In a Millikan's oil drop experiment a single negatively charged drop of radius $6 \times 10^{-6} \text{ m}$ was found to fall under gravity at a terminal velocity of 0.004 cm s^{-1} and to rise at 0.012 cm s^{-1} when a field of $2 \times 10^{-5} \text{ Vm}^{-1}$ was suitably applied. Given that the viscosity of the medium was $2.122 \times 10^{-5} \text{ Nsm}^{-2}$, determine the number of electrons on the drop.

END

NAMEPersonal NoSign			
P 510/1			
Paper 1			
DECIMALING O	e reda <i>t</i> d	TAYO EN AMINIATIONIC 2017	
BEGINNING O	F IEKWI I	TWO EXAMINATIONS, 2017	
	S.6 F	PHYSICS	
	PA	APER 1	
	2 hours	30 minutes.	
INSTRUCTIONS TO CANDIDATES:			
Attempt five questions, including at	least one ,	but not more than two from each of the sections	
A, B and C .			
Permittivity of free space ε_0	=	$8.85 \times 10^{-12} \mathrm{F m^{-1}}$	
Acceleration due to gravity, g	=	9.81 ms ⁻²	
Electronic charge, e	=	1.6 x10 ⁻¹⁹ C	
Mass of the earth	=	$5.97 \times 10^{24} \mathrm{kg}$	
Planck's constant, h	=	$6.6 \times 10^{-34} \mathrm{Js}$	
Stefan's Boltzmann's constant, σ	=	$5.7 \times 10^{-8} \text{ Wm}^{-2} \text{ K}^{-4}$	
Wien's displacement constant	=	2	
Radius of Earth's orbit about the sun	=	$1.5 \times 10^{11} \mathrm{m}$	
Radius of the sun	=	$7.0 \times 10^8 \mathrm{m}$	
Specific heat capacity of water	=	$4.2 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$	
Specific latent heat of fusion of wate	r =	$3.34 \times 10^5 \mathrm{J kg^{-1}}$	
Specific heat capacity of copper	=	$400 \text{ J kg}^{-1} \text{ K}^{-1}$	
Avogadro's number N _A	=	$6.02 \times 10^{23} \text{ mol}^{-1}$	
Density of water	=	1000 kg m^{-3}	
Gas constant, R	=	8.31 J mol ⁻¹ K ⁻¹	
Questions			
attempted			
Marks			

SECTION A

- 1(a) Write the equations of *uniformly accelerated* motion (3 marks) (b) (i) Briefly distinguish between *conservative* and *non-conservative* forces. Give two examples in each case. (4 marks) (ii) State the *principle of conservation* of energy. Show that a projectile fired with a speed $u ms^{-1}$ at an angle θ to the horizontal obeys this principle throughout its motion. (4 marks) The engine of a 5-tonne lorry can develop 30kW, and its maximum speed on level (c) road is 90 km h^{-1} . Assuming that the friction resistance is constant, calculate the greatest speed at which the lorry can climb a hill of 1 in 25. (4 marks) (d) (i)Define *moment* of a force. (1 mark) (ii) A wheel of radius 0.84 m is pivoted at its centre. A tangential force of 50 N acts on the wheel so that the wheel rotates with uniform velocity. Find the work done by the force to turn the wheel through 20 revolutions. (4 marks) 2(a) (i) Define the term angular velocity. (1mark) (ii) A car of width c and whose centre of gravity is at a height h above the ground goes round a bend of radius r. Show that it will overturn if its speed exceeds $\sqrt{\frac{c g r}{2h}}$. (5 marks) (iii) A bucket of water is swung at constant tangential speed in a vertical circle of radius 0.5 m in such a way that the bucket is upside down when it is at the top of the circle. Find the minimum speed that the bucket may have if the water is to remain in it? (3marks) (b) State Kepler's laws of gravitation. (3 marks)
 - (ii) With aid of a sketch, qualitatively explain the variation of acceleration due to gravity with distance from the centre of the earth. (3marks)

(c)

(i) Define gravitational field strength.

(1mark)

- (d) A satellite of mass m is to be launched into an orbit of radius R round the earth of radius r. Show that the minimum launching velocity is given by $v^2 = \frac{g \, r (2 \, R r)}{R}$. (4 marks)
- 3(a) (i) What is meant by the term **dimensions of a physical quantity**? (1 mark)
- (b) The volume per second of a liquid ejected from a pipe of radius a under steady flow is given by the expression $\frac{V}{t} = \frac{\pi}{8} p^x \eta^y a^z$, where p is the pressure gradient with dimensions $M L^2 T^{-2}$ and $[\eta] = M L^{-1} T^{-1}$. By dimensional analysis, find the values of x, y and z. (5 marks)
- (c) (i)Define the terms *range* and *time of flight* as applied to projectile motion.(2marks) (ii)A shell is fired from a gun with a velocity U at an angle α to the horizontal. Show that the vertical distance y covered is given by $y = x \tan \alpha \frac{gx^2 \sec^2 \alpha}{2U^2}$. (4 marks)
- (d) A shell is fired with a velocity of $100 \, ms^{-1}$ at an angle of 60° to the horizontal from a gun placed at the top of a hill of vertical height 800 m above the ground. Find the:

 (i)time taken by the shell to hit the enemy on the ground assuming the shell is on target.

 (4 marks)
 - (ii) velocity with which the shell hits the target. (4 marks)
- 4(a) (i)Distinguish between **scalar** and **vector** physical quantities. Give **two** examples of each. (4marks)
 - (ii)The diagram in figure 1 shows three coplanar forces acting on a particle.





Fig. 2

Find the magnitude, and inclination to the 8N force, of the resultant force. (5marks)

- (b) State the laws of solid friction. (3marks)
- (c) With aid of a labeled diagram, describe an experiment to determine the coefficient of static friction. (4marks)
- (d) Figure 2 shows two particles M and N each of mass 2 kg. M is held at A on an inclined plane and is connected to N by a light inelastic string that passes over a smooth pulley P. The coefficient of friction between particle M and the plane is 0.5.



Find the acceleration of the masses and the tension in the string when M is released from rest. (4marks)

SECTION B

- 5(a) (i) What is meant by a *thermometric property* and *triple point of water*? (2 marks)
 - (ii)Describe how you would measure the temperature of a body on the thermodynamic scale using a thermocouple. (3 marks)
 - (iii)The resistance of a platinum wire at the triple point of water is 5.25Ω . What will be the resistance at $100^{\circ}C$? (2 marks)
- (b) The resistance R_t of a platinum wire varies with the gas temperature t^0C according to the equation $R_t = R_0(1 + \alpha t + \beta t^2)$, where R_0 is the resistance at 0^0C and α and β

are constants. For a platinum resistance wire thermometer, α = 4.13×10⁻³ and β = -6.4×10⁻⁷ respectively. Calculate the reading of the platinum thermometer when the gas thermometer reads1000 ^{0}C . (5 marks)

- (c) Define the term **specific latent heat of vapourisation**. (1 mark)
- (d) An electrical heater rated at 500 W is immersed in a liquid of mass 2.0 kg contained in a large thermos flask of heat capacity 840 J kg⁻¹at 28°C. Electrical power is supplied to the heater for 10 minutes. If the specific heat capacity of the liquid is $2.5 \times 10^3~J~kg^{-1}~K^{-1}$, its specific latent heat of vapourization is $8.54 \times 10^3~J~kg^{-1}$ and its boiling point is $78\,^{\circ}$ C, estimate the amount of liquid which boils off.

State any assumptions made in your calculation. (7 marks)

- 6(a) (i)State **Boyle's law**. (1 mark)
 - (ii) The pressure P of an ideal gas of density ρ and mean square speed $\overline{c^2}$ is given by $P = \frac{1}{3}\rho \ \overline{c^2}$. Use this expression to derive Boyle's law. (3 marks)
- (b) Air contains approximately of 20% oxygen and 80% nitrogen. The relative molecular masses of oxygen and nitrogen are 32 and 28 respectively. Calculate the ratio of the:
 - (i)mean-square speed of oxygen to that of nitrogen. (3 marks)
 - (ii) partial pressure of oxygen to that of nitrogen in air. (4 marks)
- (c) A fixed mass of a gas is contained in a cylinder of volume V_1 and pressure P_1 . The gas is allowed to expand to a volume V_2 and pressure P_2 . Assuming that the temperature remained constant, show that the work done by the gas is given by $W = -P_1V_1\ln\frac{P_2}{P_1}$. (3 marks)
- (d) A litre of hydrogen gas at a temperature of 27°C and a pressure 10^5 Nm⁻² expands isothermally until its volume is doubled and is then adiabatically expanded until the new volume is again doubled. Given that $C_p = 29.2 \ J \ mol^{-1} \ K^{-1}$,
 - (i)Show the above processes on a P-V diagram, (2 marks)

(ii)Calculate the final pressure of the gas. (4 marks) (i) Define thermal conductivity of a material 7(a) (1 mark) (ii)Briefly account for the fact that metals are better conductors of heat than insulators. (3 marks) (b) A wall of a building consists of two brick layers each of thickness 10.0 cm and between which there is a layer of air 2.0 cm thick. Find the rate of heat flow through one m² of the wall if the inner and outer temperatures of the building are 25°C and 15°C respectively. (6 marks) (c) Describe with the aid of a labeled diagram, how the temperature of a furnace may be measured. (5 marks) The total power output of the sun is $4.0 \times 10^{26} W$. Given that the mass of the sun is (d) $1.97\times10^{30}~kg$ and its density is $1.4\times10^3~kg\,m^{-3}$, estimate the temperature of the sun. State any approximations made. (5 marks) SECTION C 8(a) Define the following terms as used in photo electricity. (i) work function. (1 mark)

(b) (i)Describe an experiment to determine Planck's constant in a laboratory. (5 marks)

(ii) stopping potential

(ii) Write down Einstein's photo-electric equation and explain how it accounts for emission of electrons from a metal surface. (3 marks)

(iii)Electromagnetic radiation of frequency 8.8×10^{14} Hz falls upon a surface whose work function is 2.5 eV. Calculate the maximum speed of the photoelectrons released from the surface. (4 marks)

(1mark)




Figure 3, shows two parallel metal plates P and Q each of length 5.0 cm and separated by a distance of 4.0 cm. A p.d. of 12V is applied between P and Q and the space between P and Q is a vacuum. A beam of electrons of speed $\,u=1.0\times10^6\,$ ms⁻¹ is directed midway between P and Q. Calculate the angle at which the beam emerges from the space between P and Q to the initial direction of the beam.

(6 marks)

- 9(a) (i) Explain how Millikan's experiment for measuring the charge of the electron proves that charge is quantized. (3 marks)
 - (ii) Oil droplets are introduced into the space between two flat horizontal plates, set 5.0 mm apart. The plate voltage is the adjusted to exactly 780V so that one of the droplets is held stationary. Then the voltage is switched off and the selected droplet is observed to fall a measured distance of 1.5 mm in 11.2 s. Given the density of oil used is 900 kgm^{-3} and the viscosity of air is $1.8 \times 10^{-5} \text{ Nsm}^{-2}$, calculate the charge of the droplet.
- (b) (i) What are cathode rays? (1 mark)
 - (ii) A horizontal beam of electrons, moving with a uniform speed, enters a uniform electric field between two horizontal parallel charged plates. Show that the path of the beam between the plates is a parabola. (3 marks)
 - (iii)An electron gun operating at $3 \times 10^3 \, V$ is used to project electrons into the space between two oppositely charged parallel plates of length $10 \, cm$ and separation $5 \, cm$. Calculate the vertical deflection of the electrons as they emerge from the region between the charged plates when the potential difference is $1.0 \times 10^3 \, V$. (4 marks)

(Use charge to mass ratio of an electron = $1.8 \times 10^{11} \text{ Ckg}^{-1}$)

(c) A beam of positive ions is accelerated through a potential difference of $1.0 \times 10^3 \text{ V}$ into a region of uniform magnetic field of flux density **0.2T**. While in the magnetic

field, it moves in a circle of radius **2.3 cm**. Derive an expression for the charge to mass ratio of the ions, and calculate its value. (4 marks)

END

S6 PHY PAPER 1, B.O.T TERM 2 EXAM 2017


Qn	Answer	
1 (a)	v = u +at	1
	$s = ut + \frac{1}{2} at$	1
	$v^2 = u^2 + 2as$	1
	where a = acceleration	
	u = initial velocity Provided the quantities	
	v = velocity after time t	
	s = displacement after time t	
(b)		
	Let the datum level be through the point of projection, O Then, at O the p.e = 0 and k.e = ½mu²	1/2
	$= \frac{1}{2} \text{m} (u_x^2 + u_y^2)$	
	$= \frac{1}{2} mu^{2} (\cos^{2}\theta + \sin^{2}\theta)$ $\therefore \text{at O the total m.e} = \text{p.e} + \text{k.e} = 0 + \frac{1}{2} mu^{2} \cos^{2}\theta + \frac{1}{2} mu^{2} \sin^{2}\theta$	1/2

	At point A, where the velocity is v at an angle α to the horizontal, the total m.e = mgh + ½mv ²	1/2
	$= \inf_{\alpha \in \mathbb{R}^n} + \frac{1}{2\pi n} \sqrt{1 + \frac{1}{2\pi n}}$ But $\sqrt{1 + \frac{1}{2\pi n}} = \sqrt{1 + \frac{1}{2\pi n}} \sqrt{1 + \frac{1}{2\pi n}}$	/2
	Now, $v^2 \cos^2 \alpha = u^2 \cos^2 \theta$ (no change in horizontal velocity)	1/2
	and $v^2 \sin^2 \alpha = u^2 \sin^2 \theta - 2gh$	1/2
	$\therefore v^2 = u^2 \cos^2 \theta + u^2 \sin^2 \theta - 2gh$	
	:. at A the total m.e = mgh + $\frac{1}{2}$ m($u^2 \cos^2 \theta + u^2 \sin^2 \theta - 2$ gh)	1/2
	$= mgh + \frac{1}{2}mu^2 - mgh$	
	= ½mu² as at point O	
(c)		
(0)		
	P P	
	$\sin\theta = \frac{1}{25}$	
	$\lambda / (0)$	
	mg sinθ At maximum speed acceleration, a = 0	
	Let P = driving force	
	Then $P - (R + mg \sin \theta) = 0$ $\therefore P = R + mg \sin \theta \dots (i)$	1
	$P = R + mg \sin\theta \dots (1)$ On the level road	_
		1
	$R = \frac{30 \times 10^3}{}$	
	V	1
	∴ R = 1200 N Let v' = maximum velocity on the slope	
		1
	Then $\frac{30 \times 10^3}{\text{v}'} = 1200 + 5 \times 10^3 \times 9.81 \times \frac{1}{25}$	
	V = 9.49 m s	
(d)	(i) The moment of a force about a point is the product of the force and the perpendicular	
(-,	distance of its line of action from the point.	1
	(ii)	
	F = 50 N; s = 0.84 m	4
	$\theta = n(2\pi) = 20 \times 2\pi = 40\pi$ radians Son Work done = Force x distance moved	1
	work done = roce x distance moved	1
	$\begin{array}{c c} 50N & = F \times \Theta S \\ & = 50 \times 40\pi \times 0.84 \end{array}$	1
	θ = 5,278 N	1
	<u> </u>	
	Total = 20	
2 (a)	(i) the rate at which a line joining a point on the body to the centre of the circle sweeps	
2 (a)	through an angle.	1
	(ii)	_
	R_1 Suppose a car of mass is moving with a	
	velocity v round a horizontal circular track	
	of radius.	
	Let R_1 and R_2 be the respective normal	
	h	1
		43
	F_1 A F_2 B	
	m -	

	reactions at the wheels A and B, and F ₁ and F ₂ the corresponding frictional forces,	1
	which provide the centripetal force.	1/2
	Then $F_1 + F_2 = \frac{mv^2}{r}$ (1) and $R_1 + R_2 = mg$ (2)	1/2
	and $R_1 + R_2 = mg$ (2) Taking moments about the centre of gravity, G, we have $ (F_1 + F_2)h + \frac{1}{2}cR_1 - \frac{1}{2}cR_2 = 0 (3) $	1/2
	Substituting for $(F_1 + F_2)$ in equation (3)	
	$\frac{mv^2}{r}h + \frac{1}{2}cR_1 - \frac{1}{2}cR_2 = 0 \qquad (4)$	
	Solving for R ₁ and R ₂ between equations (2) and (4), we get	1/2
	$R_1 = \frac{1}{2} m \left(g - \frac{2v^2 h}{cr} \right)$	1/2
	and $R_2 = \frac{1}{2} m \left(g + \frac{2v^2h}{cr} \right)$	1/2
	The car will overturn if $R_1 < 0$	
	$\therefore \frac{1}{2} m \left(g - \frac{2v^2 h}{cr} \right) < 0$	1/2
	$\therefore v > \sqrt{\frac{cgr}{2h}}$	1/2
	(iii) For the water to remain in the bucket R > 0 $ \frac{mv^2}{r} > mg $ $ \frac{mv^2}{r} > rg $ $ \frac{v^2}{r} > rg $ $ v > \sqrt{0.5 \times 9.81} $ $ v > 2.215 \text{ m s}^{-1} $	
(b)	 The planets describe ellipses about the sun as one focus. The line joining the sun and the planet sweeps out equal areas in equal times. The squares of the periods of revolution of the planets are proportional to the cubes of their 	1 1
(c)	means distances from the sun. (i) Gravitational field strength is the force acting on a mass of 1 kg placed in the gravitational	1
	field (ii) g' g (at the surface)	1/2
		1
	Inverse	44
	INSIDE OUTSIDE	
	O $x < R_E$ R_E $x > R_E$	

	T	
	~	1/2
	At points inside the earth, the acceleration due to gravity is directly proportional to the distance from the centre of the earth. At points above the earth's surface, the acceleration due to gravity varies inversely as the square	1/2
	of the distance from the centre of the earth.	1/2
(d)	Then, the energy needed, W = increase in potential energy and kinetic energy This is equal to the k.e of the satellite at launch $ \therefore \frac{1}{2} m v^2 = \frac{GMm}{r} - \frac{GMm}{R} + \frac{1}{2} m v_o^2, \text{ where } v_o \text{ is the velocity in the orbit(1)} $ But $ \frac{m v_o^2}{R} = \frac{GMm}{R^2} \implies \frac{1}{2} m v_o^2 = \frac{GMm}{2R} $ $ \therefore \frac{1}{2} m v^2 = \frac{GMm}{r} - \frac{GMm}{2R} $	
3 (a)	Dimensions of a physical quantity is the way a physical quantity is related to the fundamental quantities, i.e. mass, length and time.	1
(b)	[p] = $ML^{-2}T^{-2}$, [n] = $MT^{-1}L^{-1}$ So $\left[\frac{V}{t}\right] = \left[p\right]^{x} \cdot \left[\eta\right]^{y} \cdot \left[a\right]^{z}$ $\therefore L^{3}T^{-1} = \left(ML^{-2}T^{-2}\right)^{x} \cdot \left(MT^{-1}L^{-1}\right)^{y} \cdot L^{z}$ $\therefore L^{3}T^{-1} = M^{x+y} \cdot L^{-2x-y+z} \cdot T^{-2x-y}$ Comparing coefficients, we have For L: $-2x - y + z = 3 \dots (1)$ For M: $x + y = 0 \dots (2)$	1 1
	For ivi. $x + y = 0 \dots (2)$ For T: $-2x - y = -1 \dots (3)$ From (1) and (3): $z = 4$ Eq(2) + Eq(3): $-x = -1$ $\therefore x = 1$ And $y = -1$	1 1 1

(c)	(i) Range is the distance, measured directly, between the point of projection and the point	
(c)	where the projectile lands.	1
	Time of flight is the time between the instant of projection and landing.	1
	(ii) (x,y)	1
	Consider the bullet to be at point (x,y) on the trajectory The horizontal distance, x, after time t is given by	1/2
	$x = (u \cos \alpha)t$	1/2
	\therefore t = $\frac{x}{}$	
	u cosα	1/2
	The vertical height attained is $y = (u \sin \alpha)t - \frac{1}{2}gt^2$	1
	gx ²	
	$\therefore y = x \tan \alpha - \frac{gx^2}{2u^2 \cos^2 \alpha}$	1/2
	$\therefore y = x \tan \alpha - \frac{g x^2 \sec^2 \alpha}{2u^2}$	
(d)	$U = 100 \text{ m s}^{-1}, \ \theta = 60^{\circ}, \ h = 800 \text{ m}$	
	u Using $h = ut \sin\theta - \frac{1}{2}gt^2$	1
	$-800 = 100t \sin 60^{\circ} - \frac{1}{2} \times 9.81t^{2}$	1
	$4.905t^{2} - 50\sqrt{3} \ t - 800 = 0$ $t = 24.35 \text{ s or } -6.70 \text{ s}$	1
	∴ t = 24.35 s	1
	800m	1
	$v = \sqrt{v_x^2 + v_y^2}$	
		1
	$v_x = u \cos\theta = 100 \cos 60^\circ = 50 \text{ m s}^{-1}$ $v_y = u \sin\theta - \text{gt} = 100 \sin 60^\circ - 9.81 \text{ x } 24.35 = -152.27 \text{ m s}^{-1}$	1
	$v = \sqrt{50^2 + (-152.27)^2} = 160.27 \text{ m s}^{-1}$	1
	$v = \sqrt{50^2 + (-152.27)^2} = 160.27 \text{ m s}^{-1}$ Direction, $\alpha = \tan^{-1} \left(\frac{-152.27}{50} \right) = -71.82^{\circ}$	1
	i.e. 71.82° below the horizontal	
	(i)	
4 (a)	A scalar is a quantity which is defined by only magnitude.	1



	Applying F = ma, we have $2g - T = 2a$	½ ½
	Eq(1) + eq(2) $2g(1 - \sin 30^{\circ} - 0.5 \cos 30^{\circ}) = 4a$ $\therefore a = \frac{1}{2} \times 9.81(1 - \sin 30^{\circ} - 0.5 \cos 30^{\circ})$	1/2
	= 0.329 m s ⁻² From (1): T = 2 x 9.81 - 2 x 0.329 = 18.96 N	½ 1
	Total = 20	
5 (a)	(i) A thermometric property is a physical property of a substance whose value changes continuously with temperature. The triple point of water is a unique temperature at which pure water, pure ice and pure water vapour co-exist in equilibrium.	1 1
	(ii)	
	- The emf, E _{tr} , of the thermocouple at the triple point is measured.	1 1
	The temperature, θ , is calculated from $\theta = \frac{E_{\theta}}{E_{tr}} \times 273.16 \text{K}$	1
	(iii) $R_{tr} = 5.25\Omega$, $R_{\theta} = ?$	
	Now $373 = \frac{R_{100}}{5.25} \times 273.16$	1
	$\therefore \qquad R_{100} = \frac{373 \times 5.25}{273.16} = 7.17 \Omega$	1
(b)	$R_t = R_o(1 + \alpha t + \beta t^2), \alpha = 4.13 \times 10^{-3}, \beta = -6.4 \times 10^{-7}$	
	$t = \left(\frac{R_t - R_o}{R_{100} - R_o}\right) \times 100^{\circ} C$	1
	$= \left[\frac{R_o (1+1000 \times 4.13 \times 10^{-3} -1000^2 \times 6.4 \times 10^{-7}) - R_o}{R_o (1+100 \times 4.13 \times 10^{-3} -100^2 \times 6.4 \times 10^{-7}) - R_o} \right] \times 100^{\circ} \text{C}$	1
	-	1
	$= \frac{4.13 - 0.64}{0.413 - 0.0064} \times 100^{\circ} C$	1
	$= \frac{3.49}{0.4066} \times 100^{\circ} \text{C} = 858.34^{\circ} \text{C}$	1
(c)	This is the quantity of heat required to convert 1 kg mass of a substance from liquid to vapour at constant temperature.	1
(d)	$ \begin{pmatrix} \text{Heat supplied} \\ \text{by heater} \end{pmatrix} = \begin{pmatrix} \text{Heat gained} \\ \text{by flask} \end{pmatrix} + \begin{pmatrix} \text{Heat gained} \\ \text{by liquid} \end{pmatrix} + \begin{pmatrix} \text{Heat taken to} \\ \text{boil liquid} \end{pmatrix} $	1

	$500 \times 10 \times 60 = C_{f} \cdot \Delta\theta + m_{I}c_{I} \cdot \Delta\theta + mL_{v}$ $300,000 = 840(78 - 28) + 2 \times 2.5 \times 10^{3}(78 - 28) + m \times 8.54 \times 10^{3}$	1 3
	300,000 = 42,000 + 250,000 + 8540m	3
	m = 0.937 kg	1
	Assumptions: - All heat is taken up by the flask and the liquid.	1/2
	- No heat is lost to the surroundings	1/2
	Total = 20	
6 (a)	(i) The volume of a fixed mass of gas at constant temperature is inversely proportional to the pressure.	1
	(ii) $p = \frac{1}{3} \rho \overline{c^2}$ Let m = mass of each molecule and N = number of molecules, V = volume of gas Then $p = \frac{1}{3} \frac{Nm}{V} \overline{c^2}$	1/2
	$\therefore pV = \frac{1}{3} \text{Nmc}^{2}$ $\therefore pV = \frac{2}{3} \text{N} \left(\frac{1}{2} \text{mc}^{2} \right)$	1/2
	$ ∴pV = \frac{2}{3} N(\frac{3}{2} kT), where k = Boltzmann's constant $ $ ∴pV = NkT $ Thus, at constant temperature pV = constant	½ ½ 1
(b)	Oxygen = 20%; nitrogen = 80% $M_o = 32$, $M_N = 28$ (i) Using $p = \frac{1}{3}\rho \overline{c^2}$ $pV = \frac{1}{3}(Nm)\overline{c^2}$	
	Now, Nm = M = molecular mass	1
	For one mole of oxygen $p_oV = \frac{1}{3}M_o c_o^2 = RT$	1
	and for one mole of oxygen $p_n V = \frac{1}{3} M_n \overline{c_n^2} = RT$	_
	$\frac{\therefore \text{ at the same temperature,}}{\overline{c_o^2} : \overline{c_n^2}} = \frac{3RT}{32} : \frac{3RT}{28} = 28 : 32 = 7 : 8$	1
	(ii) Let P _o = partial pressure due to oxygen P _n = partial pressure due to nitrogen n _o = number of oxygen molecules in the mixture n _n = number of nitrogen molecules in the mixture m _o = mass of each oxygen molecule m _n = mass of each nitrogen molecule	
	Then $P_o = \frac{1}{3} \frac{n_o m_o}{V} \overline{c_o^2}$ and $P_n = \frac{1}{3} \frac{n_n m_n}{V} \overline{c_n^2}$	1
	$\therefore \frac{P_{o}}{P_{n}} = \frac{n_{o} m_{o}}{n_{n} m_{n}} \cdot \frac{\overline{c_{o}^{2}}}{\overline{c_{n}^{2}}} = \frac{20}{80} \cdot \frac{7}{8} = \frac{7}{32}$	2

(c)	The total world done in companding from M. to M. is		1
(c)	The total work done in expanding from V_1 to V_2 is		
	$\mathbf{P} = \int d\mathbf{W}$	$V = \int_{V}^{V_2} P dV$	1
		V ₁	
	For n moles of nRT	rgas	
	$P = \frac{nRT}{V}$		
	V_1 V_2 V	AV.	1
	Hence $W = \int_{V_1}^{V_2} PdV = \int_{V_1}^{V_2} nRTdV = nRTlog$	$v_e\left(\frac{v_2}{V_1}\right)$	
	The heat required, $Q = W = nRTlog_e \left(\frac{V_2}{V_1}\right)$		
	But $P_1V_1 = P_2V_2 = nRT$		
	$\therefore V_2 = \frac{P_1 V_1}{P_2}$		
	2	(D) X	1
	$\therefore W = P_1 V_1 \log_e \left(\frac{P_1 V_1}{P_2} \cdot \frac{1}{V_1} \right) = P_1 V_1 \log_e \left(\frac{P_1}{P_2} \right) = -P_1 V_1 \log_e \left(\frac{P_1}{P_2} \right)$	$=\left(\frac{\mathbf{r}_2}{\mathbf{P}_1}\right)$	
(d)	$V_1 = 1 \text{ litre} = 1 \times 10^{-3} \text{m}^3$ $T_1 = 27^{\circ}\text{C} = 300 \text{ K}$ $P_1 = 10^5 \text{ N}$	m ⁻²	
	$V_2 = 2 \times 10^{-3} \text{ m}^3$ $V_2 = T_1$ $P_2 = ?$ $V_3 = 4 \times 10^{-3} \text{ m}^3$ $T_3 = ?$ $P_3 = ?$		
	3		
			1/2
	$P_1 \longrightarrow A$ AB = Iso	thermal	
	BC = Ar	diahatic	1/2
	P ₂		
	P ₂		1/2
	P ₃ C		1/
	V_1 V_2 V_3 V_3		1/2
	Along AB: $P_1V_1 = P_2V_2$		1/2
	$\therefore P_2 = \frac{10^5 \times 1 \times 10^{-3}}{2 \times 10^{-3}} = 5 \times 10^4 \text{ Nm}^{-2}$	$C_p = 29.2 \text{ J mol}^{-1} \text{ K}^{-1}$	1/2
	$\left(2 \times 10^{-3}\right)^{1.40}$	$C_v = C_p - R$ = 29.2 - 8.83	1/2
	$P_3 = 5 \times 10^4 \left(\frac{2 \times 10^{-3}}{4 \times 10^{-3}} \right)^{1.40}$	= 29.2 – 8.83 = 20.89 J mol ⁻¹ K ⁻¹	1/2
	= 18,950 Nm ⁻²	= 20.89 J mol ⁻¹ K ⁻¹	1
	Using $T_2V_2^{\gamma-1}=T_3V_3^{\gamma-1}$	I	

	$T_3 = 300 \left(\frac{2 \times 10^{-3}}{4 \times 10^{-3}} \right)^{1.40-1} = 227.4 \text{ K}$	1½
	Total = 20	
7 (a)	(i) Thermal conductivity is the heat flow rate in a substance per unit area per unit temperature gradient.	1
	 (ii) In poor conductors heat is conducted by waves produced by lattice vibrations. This happens, say, when heat is applied to one part of the solid, the kinetic energy of vibration of the molecules there is increased. Because of the bonds between the molecules, the kinetic energy of vibration of the neighbouring molecules is in turn also increased. So the wave moves on. In metals conduction is predominantly due to freely moving electrons (in addition to lattice vibration). When a part of a metal is heated, the free electrons gain thermal energy and their velocities 	y ₂ y ₂ y ₂
	increase.They distribute this energy by collision with positive ions in the lattice and increase the ions' vibrational energy.	½ ½
	- Because electrons are light, they are able to move quickly to the cooler parts of the solid. So this mode of heat transfer is much faster.	1/2
(b)	Assume the heat flow rate through the layers to be same. $ 25^{\circ}C \qquad \theta_1 \qquad \theta_2 \qquad 15^{\circ}C \qquad \checkmark \\ \qquad \qquad k_b k_a \qquad k_b \qquad \qquad k_b $	1/2
	$\frac{k_b A(25 - \theta_1)}{0.1} = \frac{k_a A(\theta_1 - \theta_2)}{0.02} = \frac{k_b A(\theta_2 - 15)}{0.1}$	1 1/2
	$\therefore \frac{0.7(25 - \theta_1)}{0.1} = \frac{0.024(\theta_1 - \theta_2)}{0.02} = \frac{0.7(\theta_2 - 15)}{0.1}$ $\therefore \frac{0.7(25 - \theta_1)}{0.1} = \frac{0.7(\theta_2 - 15)}{0.1}$	
	$\therefore 25 - \theta_1 = \theta_2 - 15$ $\therefore \theta_1 + \theta_2 = 40 \dots (1)$	1/2
	Also $7(25 - \theta_1) = 1.2(\theta - \theta_2)$ $\therefore 8.2\theta_1 - 1.2\theta_2 = 175$ (2) From (1): $\theta_1 = 40 - \theta_2$ Substituting for θ_1 into (2):	1/2
	Substituting for θ_1 into (2): $8.2(40 - \theta_2) - 1.2\theta_2 = 175$	½ ½

	$\therefore \qquad 9.4\theta_2 = 153$ $\therefore \theta_2 = 16.3^{\circ}C$	1
	1 -	1
	:. Heat flow rate per $m^2 \frac{0.7(16.3-15)}{0.1} = 9.1 \text{Wm}^{-2}$	
(c)	A pyrometer is used. What is shown below is an optical pyrometer	
	O Filament E Correct labeling of any 4 main parts @1/2	2
	 It consists of a telescope, OE, and a lamp having a tungsten filament. G is a red filter through which light from the body, B, whose temperature is required passes. The eyepiece, E, is focused upon the filament. The furnace, B, is then focused by the objective lens O so that its image lies in the plane of the filament. The temperature of the filament is adjusted using rheostat R until it "disappears" in the background of the radiation from B. Now, the ammeter, A, which measures the current, has been calibrated directly in degrees, and gives the temperature of the furnace. 	
(d)	Power output = 4.0×10^{26} W m _s = 1.97×10^{30} ; ρ_s = 1.4×10^3 kg m ⁻³ Assuming that the sun is spherical	
	Volume, V = $\frac{\mathrm{m}}{\mathrm{m}} = \frac{4}{3} \pi \mathrm{r}_{\mathrm{s}}^3$	
	$\therefore 3 \times \frac{1.97 \times 10^{30}}{1.4 \times 10^{3}} = 4\pi r_{s}^{3}$	1
	$\therefore r_s = \sqrt[3]{3.359 \times 10^{26}} = 6.952 \times 10^8 \text{m}$	1
	Assume that the sun radiates as a black body	
	Then power radiated by the sun = $4.0 \times 10^{26} \text{ W} = \sigma \text{A} \text{ T}_s^4$	
	$\therefore 4\pi r_s^2 \sigma T_s^4 = 4 \times 10^{26}$	1
	$T_s^4 = \frac{4 \times 10^{26}}{4\pi \times (6.952 \times 10^8)^2 \times 5.67 \times 10^{-8}}$	1
	$\therefore T_{s} = \sqrt[4]{1.162 \times 10^{15}}$	
	= 5,838 K	1
Total = 20		

P 510/1 PHYSICS
Paper 1
UGANDA ADVANCED CERTIFICATE OF EXAMINATIONS
PHYSICS
S.6 PAPER 1 - 2014
2 hours 30 minutes.
INSTRUCTIONS TO CANDIDATES:
Attempt five questions in all.
Non-programmable scientific electronic calculators may be used.
Assume where necessary.

Acceleration due to gravity, g 9.81 ms⁻² $5.97 \times 10^{24} \text{ kg}$ Mass of the earth = $5.7x10^{\text{-8}}\;Wm^{\text{-2}}\;K^{\text{-4}}$ Stefan's Boltzmann's constant, σ Radius of the earth $6.4 \times 10^{6} \text{m}$ Radius of the sun $7.0 \times 10^8 \text{m}$ Radius of Earth's orbit about the sun $1.5 \text{ x} 10^{11} \text{m}$ Speed of light in a vacuum, *c* $3.0 \times 10^{8} \text{ms}^{-1}$ Specific heat capacity of water 4,200J kg⁻¹ K⁻¹ Universal gravitational constant, G $6.67 \ x10^{-11} N \ m^2 \ kg^{-2}$ 1000 kgm⁻³ Density of water 8.31 J mol⁻¹K⁻¹ Gas constant, R **SECTION A** 1. (a) State Kepler's laws of planetary motion. (3) (b) Derive the expression for the centripetal force on mass \mathbf{m} moving with uniform speed v in a circular path of radius r. (4) (c) Explain why: (i) the earth has atmosphere. (2)

- (ii) a cyclist has to lean over towards the centre of a bend in order to turn well. (2)
- (d) A satellite of mass 120 kg moves in a circular orbit around the earth with a period of 7.7×10^7 s.
- (i) Find its height above the earth. (5)
- (ii) Calculate the mechanical energy of the satellite. (4)
- 2. (a) What is meant by *simple harmonic motion*? (1)
 - (b) A cylindrical vessel of cross-sectional area **A** contains air of volume **V**, at pressure **P**, trapped by a frictionless piston of mass **M**. The piston is pushed down and then released.
 - (i) If the piston oscillates with simple harmonic motion, show that its frequency, f, is given by

$$f = \frac{A}{2\pi} \sqrt{\frac{P}{MV}} \tag{6}$$

- (ii) Show that the expression for f, in b(i) is dimensionally correct. (2)
- (c) A mass of **0.1kg** suspended from a spring of force constant **24.5 Nm**⁻¹ is pulled vertically downwards through a distance of **5.0 cm** and released. Find the

	(i) period of oscillation.	(2)
	(ii) position of the mass 0.3 seconds after release.	(4)
	(d) Sketch the following graphs for a body performing simple harmonic motion:	
	(i) Velocity against displacement	(1)
	(ii) Displacement against time.	(1)
	(e) With the aid of sketch graphs, distinguish between free and damped oscillation	s. (3)
3.	(a) (i) Distinguish between <i>kinetic energy</i> and <i>potential energy</i> and show that a mam, moving with a velocity, v, has kinetic energy given by $^1\!/_2mv^2$	ss (5)
	(ii) What is meant by the term <i>conservative force</i> ?	(1)
	(iii) Show that for a particle moving in the gravitational field the total mechanical energy is conserved.	(4)
	(b) (i) Define the <i>moment of a force</i> .	(1)

	(ii) The inner end of a concentrically coiled spring is fixed to the axle of a wheel o radius 0.5 m . When two tangential parallel forces, each of 6N acting in opposite directions, are applied to the wheel to form a couple, the wheel turns through an angle of 120° . Find the energy stored in the spring.	f (3)
	(c) The engine of a 5-tonne lorry can develop 30kW , and its maximum speed on loroad is 90 km h ⁻¹ . Assuming that the friction resistance is constant, calculate the greatest speed at which the lorry can climb a hill of 1 in 25 .	evel (4)
	(d) Explain rocket propulsion.	(3)
	SECTION B	
4.	(a) (i)Explain why the difference between the principal heat capacities of a solid in egligible.	s (2)
	(ii) Derive an expression for the difference in molar heat capacities of a gas.	(4)
	(b) (i) Explain why the sound propagation in air is considered to be an adiabatic process.	(2)
	(ii) A fixed mass of gas at a pressure P_1 and volume V_1 expands isothermally to a pressure P_2 and volume V_2 . Derive an expression for the work done by the gas.	(4)

(c) A gas of volume 2 litres at a temperature of 27 °C and pressure of 1.5 x 10 5 Pa is	S
heated at constant pressure until its volume doubles. It is then cooled at constant	
volume back to its original temperature before finally being compressed	
isothermally to its original volume.	
Draw a p-V diagram of the whole cycle and find the net work done by the gas.	(5)
(d) The pressure P of an ideal gas of density ρ , is given by $P = \frac{1}{3} \rho \ \overline{c^2}$, where $c^{\overline{2}}$ is	
the mean-square speed of its molecules.	's
hypothesis.	(3)
(a) Explain why	
(i) at low pressures a real gas approximates to a perfect gas.	(2)
	(2)
(ii) evaporation causes cooling.	(2)

5.

cylinder and explain its features. (3)

(b) (i) State two properties of a saturated vapour.

(iii) Describe an experiment to show that saturated vapour pressure varies with temperature. (6)

(2)

	(c) Air saturated with water vapour and contained in a rigid cylinder, at 100°C are	ıd
	2.0 x 10 ⁵ Pa pressure is cooled to 20°C . Neglecting the expansivity of the cylinde calculate the final pressure in the cylinder, if the saturated vapour pressure of wa	
	20 °C is 2.3 x 10 °3 Pa. [Atmospheric pressure = 1.013 x 10 °5 Pa]	(5)
6.	(a) Define thermal conductivity of a material.	(1)
	(b) (i) Using molecular theory of matter, explain the mechanism of heat energy transfer in insulators.	(3)
	(ii) Briefly account for the fact that metals are better conductors of heat than insulators.	(3)
	(c) A pan of diameter 20.0 cm and thickness 2.0 mm is filled with water and heat such that the water boils producing steam at a rate of 5.0g s ⁻¹ at steady state conditions.	ed
	(i) calculate the temperature of the external surface of the pan.	(4)
	(ii) State the assumptions made.	(2)
	(Specific latent heat of vaporisation of water is $2.26 \times 10^6 J \text{ kg}^{-1}$ while conductivit the material of the pan is $380 \text{ Wm}^{-1} \text{ K}^{-1}$)	y of

(d) (i) What is meant by black body radiation? (1)

(ii) The total power output of the sun is 4.0×10^{26} W. Given that the mass of the sun is 1.97×10^{30} kg and its density is 1.4×10^{3} kg m⁻³, estimate the temperature of the sun. State any assumptions made.

(Stefan's constant,
$$\sigma = 5.67 \times 10^{-8} \text{ Wm}^{-1}\text{K}^{-1}$$
) (6)

END

S6 PHYSICS

Paper 1

2 ½ hours

NAME		n H'se
Signature		
Answer ALL questions.		
Acceleration due to gravity, g	=	9.81 ms ⁻²
Mass of the earth	=	$5.97 \times 10^{24} \text{ kg}$
Stefan's Boltzmann's constant, σ	=	5.67x10 ⁻⁸ Wm ⁻² K ⁻⁴
Radius of the earth	=	6.4 x 10 ⁶ m
Radius of the sun	=	$7.0 \times 10^{8} \text{m}$

Radius of Earth's orbit about the sun = $1.5 \times 10^{11} \text{m}$

Specific heat capacity of water = $4,200 \text{J kg}^{-1} \text{ K}^{-1}$

Universal gravitational constant, G = 6.67 x10⁻¹¹N m⁻² kg⁻²

Density of water = 1000 kgm^{-3}

Gas constant, R = 8.31 J mol⁻¹K⁻¹

Question	1	2	3	4	5	TOTAL
Marks						

SECTION A

1(a) (i) State the principle of conservation of mechanical energy. (01)

- (ii) A particle is projected at an acute angle, **\(\theta\)**, to the horizontal. Assuming no air resistance, show that the mechanical energy of the particle is conserved throughout its motion. (04)
- (b) A ball of mass **0.5 kg** is allowed to drop from rest, from a point a distance of **5.0 m** above a horizontal concrete floor. When the ball first hits the floor, it rebounds to a height of **3.0m**.
 - (i) What is the *speed of the ball just after the first collision* with the floor? (04)
- (ii) If the collision lasts 0.01 s, find the *average force* which the floor exerts on the ball. (04)
- (c) A bullet of mass **10 g** is fired at short range into a block of wood of mass **990 g** resting on a smooth horizontal surface and attached to a spring of force constant **100 Nm**-1. The bullet remains embedded in the block while the

(d)	Explain, giving two examples, the meaning of a 'conservative force'	(02)
2.(a)	(i) State the <i>laws of solid friction</i> .	(03)
	(ii) Explain, using the molecular theory, the laws stated in (i) above.	(03)
(b)	Describe how the <i>coefficient of static friction</i> for an interface between rectangular block of wood and a plane surface can be determined.	
(c) coplai	State the <i>conditions for the equilibrium</i> of a rigid body under the actual forces. (02)	
(d)	A mass of 5.0 kg is suspended from the end A of a uniform beam of kg and length 1.0 m . The end B of the beam is hinged in the beam is kept horizontal by a rope attached to the mid-point of the bar a point C , in the wall at a height 0.75 m above B . Calculate:	wall. The
	(i) the <i>tension</i> in the rope.	(05)
	(ii) the force exerted by the hinge on the beam.	(03)
3.(a)	(i) Define angular velocity.	(01)

(ii) Derive an expression for the force F, on a particle of mass m, moving with


angular velocity $\boldsymbol{\omega}$, in a circle of radius \boldsymbol{r} .

spring is compressed by a distance of **5.0 cm**. Find the *elastic potential energy* of the compressed spring, and the *speed of the bullet* just before collision with

(05)

the block.

(04)



A pendulum bob is set rotating so that it describes a horizontal circle with angular velocity ω as shown in the figure above. If the mass of the bob is \mathbf{m} and the length of the string is \mathbf{d} , find in terms of \mathbf{m} , \mathbf{d} the expression for:

- (i) the tension in the string in terms of \mathbf{m} , \mathbf{d} and $\boldsymbol{\omega}$ (03)
- (ii) the tangential velocity of the bob. (03)
- (c) A body of mass, **m**, is attached to the end of a string of length, **a**. The other end of the string is fixed. The string is initially held taut and horizontal and the body is then released.

Find the velocity of the body and the tension in the string when the string reaches the vertical position. (03)

(d) (i) Derive an expression for the *speed* with which a car can negotiate a bend on *a banked track* without skidding. (03)

(ii) Explain why a <i>racing car can travel faster</i> on a banked track of the same radius of curvature.	track than (03)	on a flat
SECTION B.		
4. (a) (i) Define the two principal molar heat capacities of a gas.		(02)
(ii) State, which one is greater, and explain why this is so		(02)
(iii) Show that the difference between the molar heat cape molar gas constant.	<i>acities</i> is equ 05)	ıal to the
(b) State the conditions for an adiabatic process.		(02)
(c) A gas at a pressure of 1.2 x 10⁶ Pa and temperature 90 adiabatically to twice its original volume and then is comp to its original volume. [Take ratio of the principal molar heat cap	ressed is	othermally
(i) Sketch and label the two stages on a P-V diagrar	n.	(02)
(ii) Find the <i>final pressure</i> and <i>temperature</i> of the g	gas.	(07)

5 (a)	(i) Define thermal conductivity.	(01)
	(ii) Explain the mechanism of heat conduction in metals.	(03)
(06)	(iii) Describe an experiment to determine the thermal conductivity of	a metal.
(b)	(i) What is a <i>black body</i> ?	(01)
radiat	(ii) Draw sketch graphs to show the variation of relative intensity of b tion with wavelength for different temperatures. (02)	lack body
tempe meltin		nen its elow
	The heat radiation <i>per second per m</i> ² received by the earth from the Wm -2. Assuming this is 90 % of what the sun emits as a black be attentionate the temperature of the sun. (04)	

END

S.6 PAPER 1 TEST – (Mechanics and Modern)

1(a)	(i) Define surface tension in terms of surface energy.	(1)
	(ii) Calculate the work done against surface tension forces in blowing a soap diameter 15 mm. Take the surface tension of the soap solution as 3.0×10^{-2}	
	(iii) Lycopodium powder is sprinkled over the surface of water in a dish. A st of soap solution is then introduced at the centre of the surface.	mall drop
	Explain what is observed on the surface.	(3)
(b)	Describe an experiment to investigate the effect of temperature on surface tension of a liquid.	(6)
(c)	(i) Derive an expression for the capillary rise of a liquid of density ρ , surface and angle of contact θ , in a tube of radius, r.	tension y, (3)
	(ii) The internal diameter of the tube of a mercury barometer is 3.00mm. First corrected reading of the barometer after allowing for the error due to surfact if the observed reading is 76.56 cm. [Surface tension of mercury = 4.80×10^{-6} angle of contact of mercury with glass is 140^{-6} ; density of mercury is 13600×10^{-6}	te tension, 1 Nm ⁻¹ ;
2(a)	What is meant by the following terms:	
	(i) nuclear number.	(1)
	(ii) binding energy.	(1)

(b) Calculate the energy released by 1.0 g of a sample during the decay of $\frac{220}{86}$ Rn nucleus into $\frac{216}{84}$ Po nucleus and an alpha particle. (4)

Mass of 220 Rn = 219.964176 u

Mass of 216 Po = 215.955794 u

Mass of 4 He = 4.001566 u

1 u = 931 MeV and Avogadro's number = 6.02×10^{23}

(c) (i)With the aid of a labeled diagram, describe how the diffusion cloud chamber can be used to detect ionizing radiation from a radioactive source.

(5)

- (ii) Draw and explain the nature of tracks expected in c(i) above if the source emits α -particles. (2)
- (d) (i) Explain how you would use the decay curve for a radioactive material to determine its half life. (2)
 - (ii) A radioactive source contains 1.0 μg of plutonium of mass number 239. If the source emits 2300 alpha particles per second, calculate the half-life of Plutonium.

[Assume the decay law N = N₀ $e^{-\lambda t}$] (5)

S.6 PHYSICS: MECHANICS & HEAT

1 (a) (i) Define pressure.	(1)	
(ii) Derive the expression for the pressure at a point a depth, ${\boldsymbol h}$ from the surface of a liqu ${\boldsymbol \rho}$.	id of density,	
(b) A block of mass 0.10 kg is suspended from a spring balance. When the block is immed of density 1000 kgm -3, the spring balance reads 0.63 N . When the block is immersed in a unknown density, the spring reads 0.70 N . Find:		
(i) the density of the solid.	(4)	
(ii) the density of the liquid.	(3)	
(c)(i) State Bernoulli's principle.	(1)	
(i)Explain qualitatively the origin of the lift force on an aero-plane at take off.	(3)	
(ii)An aero-plane has a wing area of 40 m² . At take off, the speed of air above and below the wings are 120 ms⁻¹ and 100 ms⁻¹ respectively. Find the lift on the aero-plane, if the density of air is 1.3 kgm⁻³ . (5)		
2. (a) Derive the pressure $P = \frac{1}{2} \rho c^2$ for the pressure, P , of an ideal gas of density ρ is square speed c^2 . State any assumptions made.	and mean	

. , 0	of volume 0.1 m^3 at a pressure of $1.0 \times 10^5 \text{ Nm}^{-2}$ and a temperature ideal, calculate the density of the gas (The relative molecular (5)
(c) Explain how cooking at a press achieved on top of high mountain	are of 76 cm of mercury and a temperature of 100 °C , may be . (3)
•	rygen atoms are 1.66 x 10 ⁻²⁷ kg and 2.66 x 10 ⁻²⁶ kg . What is the rati Hydrogen to that of Oxygen molecules at the same temperature?
(e)Indicate the different states of a sketch graph.	real gas at different temperatures on a pressure versus volume (2) END
S.	5 PAPER 1 TEST – October, 2014
1(a) (i) State the difference	petween isothermal and adiabatic expansion of a gas. (2
	ans point, sketch graphs of pressure versus volume for a fixed isothermal and adiabatic changes.
(b) Show that the work dor given by $\int_{V_0}^{V_1} P \ dV$.	e, W , by a gas which expands reversibly from V_0 to V_1 is (4

(i) State the differences between real and ideal gases.

(c)

(2)

	(ii) Draw a labeled diagram showing P – V isothermals for a real gas above and bel the critical temperature.	ow (3)
(d)	Ten moles of a gas, initially at 27° C are heated at constant pressure of 1.01×10^{5} Pa and the volume increased from 250 litres to 375 litres. Calculate the increase in internal energy. [Assume $C_P = 28.5 \text{ Jmol}^{-1}\text{K}^{-1}$, $R = 8.314 \text{ Jmol}^{-1}\text{K}^{-1}$]	(6)
2(a)	(i) What is meant by thermionic emission?	(1)
	(ii) Sketch the Current – Potential difference characteristics of a thermionic diode two different operating temperatures and explain their main features.	for (5)
	(iii) Describe one application of a diode.	(2)
(b)	(i) What features of an X-ray tube make it suitable for continuous production of X rays?	(3)
	(ii) Sketch a graph of intensity versus frequency of radiation produced in an X-ray tube and explain its features.	(5)
(c)	A monochromatic X-ray beam of wavelength $1.0 \times 10^{-10} \text{m}$ is incident on a set of planes in a crystal of spacing $2.8 \times 10^{-10} \text{m}$. What is the maximum order possible withese X-rays?	th (4)

3(a)	Define the following terms:	
	(i) centripetal acceleration.(ii) angular velocity.	(1) (1)
(b)	(i) What is meant by banking of a track?	(1)
	(ii) Derive an expression for the angle of banking, θ , for a car of mass, m, moving a speed, v, round a banked track of radius, r.	t a (4)
(c)	A bob of mass, m, is tied to an inelastic thread of length, l, and whirled with const speed in a vertical circle.	ant
	(i) With the aid of a sketch diagram, explain the variation of tension in the string along the circle.	(5)
	(ii) If the string breaks at one point along the circle, state the most likely position explain the subsequent motion of the bob.	and (2)
(d)	A body of mass 15 kg is moved from the earth's surface to a point 1.8×10^6 m above the earth. If the radius of the earth is 6.4×10^6 m and its mass is 6.0×10^{24} kg, calcuthe work done in taking the body to that point. [use $G = 6.67 \times 10^{-11}$ Nm ² kg ⁻²]	

END

Instruction: Attempt at least two questions in one hour.

P510/1 PHYSICS
Paper 1
AUGUST 2015
UACE MOCKS 2015
PHYSICS
PAPER 1
2 hours 30 minutes.
INSTRUCTIONS TO CANDIDATES:
Attempt five questions, including at least one , but not more than two from each of the sections A, B, and C .
Assume where necessary.

Permittivity of free space ε_0 = 8.85 x 10⁻¹² Fm⁻¹

Acceleration due to gravity, $g = 9.8 \text{ ms}^{-2}$

Electron charge, e = 1.6 x10⁻¹⁹ C

Stefan's Boltzmann's constant, $\sigma = 5.7x10^{-8} \text{ Wm}^{-2} \text{ K}^{-4}$

Radius of the earth = 6.4×10^6 m

Radius of the sun = $7.0 \times 10^8 \text{m}$

Radius of Earth's orbit about the sun = 1.5×10^{11} m

Speed of light in a vacuum, $c = 3.0 \times 10^8 \text{ms}^{-1}$

Specific heat capacity of water = $4,200 \text{J kg}^{-1} \text{ K}^{-1}$

Universal gravitational constant, G = $6.67 \times 10^{-11} \text{N m}^{-2} \text{ kg}^{-2}$

Avogadro's number N_A = $6.02 \times 10^{23} \text{ mol}^{-1}$

Density of water = 1000 kgm^{-3}

Gas constant , R = $8.31 \,\mathrm{J} \,\mathrm{mol}^{-1} \mathrm{K}^{-1}$

Charge to mass ratio, $^{e}/m$ = 1.8 x 10 11 C kg $^{-1}$

The constant . $1 = 9.0 \times 10^9 \,\text{F}^{-1}\text{m}$

 $4\pi\epsilon_{o}$

SECTION A

1. (a) (i) Define the coefficient of surface tension.

(1)


(ii) Lycopodium powder is sprinkled over the surface of water is soap solution is then introduced at the centre of the surface.	n a dish. A small drop of
Explain what is observed on the surface.	(3)
(b) Describe an experiment to investigate the effect of temperature of a liquid.	e on surface tension (6)
(c) (i) State Bernoulli's principle.	(1)
(ii) Explain how the principle is applied in a filter pump.	(3)
(d) Water flows steadily along a uniform tube of cross-section 30 cr pressure is 1.25×10^5 Pa and the total pressure is 1.30×10^5 Pa	
(i) the flow velocity	(4)
(ii) the mass flow rate.	(2)
2. (a) Define the following	
(i) a couple	(1)
(ii) work	(1)

(b) Show that when a couple rotates a body the work done is equal to the product of the

torque and the angle in radians turned through.

(4)

(c) The diagram below shows three coplanar forces acting on a particle.




Find the magnitude, and inclination to the 12N force, of an additional force that will bring about equilibrium. (6)

(d) The figure below shows a uniform rigid plank AB of weight 100 N and length, 5 m freely hinged at end B in a vertical wall. A light rope is fixed between end A and a point C 5 m vertically above B so that AB makes an angle of 30° with the horizontal. A box of weight 200 N is hung from point D on the plank 2 m from end B. Find

(i) the tension in the rope. (3)

(ii) the reaction at the hinge.

(5)



В


3. (a) (i) Derive an expression for the tangential speed of a particle describing a circle of radius r at an angular velocity ω . (3)

(ii) A coin is placed on a horizontal turntable at a point 5 cm from the centre of the turntable. If the coefficient of friction between the coin and the turntable is 0.5, find the maximum frequency of the turntable for the coin to remain at the same point.(3)

(b) (i) Define gravitational potential.

(1)

- (ii) Find an expression for the total energy of a satellite of mass m circling the earth of mass M in an orbit of radius R. (3)
- (iii) Explain the effect of friction between such a satellite and the atmosphere in which it moves. (4)
 - (c) (i) Define simple harmonic motion. (1)
- (ii) The figure below shows two springs of respective constants k_1 and k_2 joined in series. A mass m resting on a smooth horizontal surface is fixed to one end of the combination.



The other end of the combination is fixed so that the springs are horizontal. If m is now pulled a short distance along the surface and then released, show that it executes simple harmonic motion and find the period. (5)

- 4. (a) (i) What is meant by a conservative force? (1)
 - (ii) Show that if a body is moving in the gravitational field the mechanical energy is conserved. (4)

(b) A light helical spring gives a displacement of 4 cm for a load of 400 supports a light pan. A mass of 100 g is dropped from a height of 4 cm or				
(i) the maximum displacement produced	(4)			
(ii) the speed with which the pan-mass combination returns to the i the pan.	nitial position of (3)			
(c) (i) What is meant by uniform acceleration?	(1)			
(ii) A particle is projected vertically upwards from the top of a tree. Stime and a velocity-time graph for the motion up to the time the particle(2)	•			
(d) A car, which is slowing down steadily, passes a point A with a speed instant a motorcyclist at A is starting off in the same direction. If the maintains a steady acceleration of 2.5 ms ⁻² and catches up with the the car's speed is 18 km h ⁻¹ , find	of 72 km h ⁻¹ at the motorcyclist car at point B when			
(i) the time taken to move from A to B	(3)			
(ii) the distance AB	(2)			
SECTION B				
5 (a) (i) Define thermal conductivity	(1)			
(ii) Explain the mechanism of heat transfer in metals	(3)			

(iii) Describe an experiment to determine the thermal conductivity of a me	etal. (6)	
(b) (i) What is a black body?	(1)	
(ii) Draw sketch graphs to show the variation of relative intensity of black be radiation with wavelength for different temperatures.	oody (2)	
(iii) Explain the appearance of a metal ball placed in a dark room when its is progressively raised from room temperature to just below melting. (3)	tempe	rature
(c) A heating element in form of a cylinder 30 cm long and 1.5 cm in diameter of 1.8 kW. If its radiation is 85% that of a black body, find its temperature. (4)	er has an	output
6. (a) Explain why (i) at low pressures a real gas approximates to a perfect gas.	(2)	
(ii) evaporation causes cooling.	(2)	
(b) (i) State two properties of a saturated vapour.(ii) Air is trapped between a piston and a liquid in a cylinder. The piston is	(2) slowly	pulled
outwards at constant temperature and this continues even after all the liquid		

has	eva	por	ate	d.

Sketch a graph of pressure against volume for the contents of the cylinder and explain its features. (3)

- (iii) Describe an experiment to show that saturated vapour pressure varies with temperature. (6)
- (c) Air saturated with water vapour and contained in a rigid cylinder, at 100°C and

 2.0×10^5 Pa pressure, is cooled to 20° C. Neglecting the expansivity of the cylinder calculate the final pressure in the cylinder, if the saturated vapour pressure of water at 20° C is 2.3×10^3 Pa.

[Atmospheric pressure =
$$1.013 \times 10^5 \text{ Pa}$$
] (5)

- 7. (a) (i) Define specific latent heat of fusion.
 - (ii) Describe an experiment to determine the specific latent heat of fusion of water (5)
- (b) (i) Explain why the molar heat capacity of an ideal gas at constant pressure differs from that at constant volume. (3)
 - (ii) Derive an expression for the difference of the above molar heat capacities. (4)
 - (c) (i) State the conditions necessary for a reversible adiabatic process. (2)
 - (ii) A gas initially occupying a volume of 1.0 litre at a temperature of 47° C and a pressure of 1.40×10^{5} Pa expands isothermally to a volume of 2.0 litres. It is then

(1)


compressed adiabatically to its original volume. If γ = 1.4 find the final temperature and pressure. (5)

SECTION C		
8. (a) (i) Distinguish between X-rays and cathode rays.	(2)	
(ii) In an X-ray tube explain the features adopted for the structure and anode. (4)	materials of	the
(b) (i) State Bragg's law.	(1)	
(ii) What is the condition for obtaining many orders of X-ray diffraction	n. (1)	
(iii) A monochromatic beam of X-rays of wavelength 1.187×10^{-10} m is in parallel atomic planes of spacing 3.00×10^{-10} m. Determine the maximum of (3)		a set of
(c) (i)In photo-electricity, what is meant by the work function?	(1)	
(ii) Describe how you could determine Planck's constant in a school lab	oratory. (5)	
(iii) When monochromatic light of frequency 6.0×10^{14} Hz falls on a met stopping potential is 0.6 V while when the same surface is struck by light of		the

frequency 1.0 x 10^{15} Hz the stopping potential becomes 2.2 V. Determine the work function of the metal. (3) 9. (a) (i) Distinguish between radioactivity and nuclear fission. (2) (ii) Why are neutrons preferred to charged particles for inducing nuclear reactions. (2) (b) Describe how a Geiger-Muller tube works. (6) (c) 1.2g of a substance forms a point source of γ -rays. Only 1 in 10¹² of its atoms are radioactive and the half-life is 100 days. A Geiger-Muller tube facing the source at a distance of 10 cm gives a count rate of 11 s⁻¹. If the window of the tube has an cm² find (i) the number of radioactive atoms present (5) (ii) the mass number of the substance (2) (d) Determine whether the nucleus ²¹⁰Po is stable or it may undergo disintegration to produce $^{206}_{82}Pb$ and an $\alpha\text{-particle}.$ (3) Mass of 210 Po = 209.937u Mass of 206 Pb = 205.929u

Mass of ${}^{4}He = 4.002u$

- 10. (a) (i) Sketch the I-V characteristic for gaseous conduction. (2)
 - (ii) Explain the main feature of the curve. (3)
 - (b) A valve diode D was connected to rectify a.c supply to a load as shown below.C is a capacitor.



- (i) State the purpose served by the capacitor C. (1)
- (ii) Explain the effect on the p.d's V_d and V_L of introducing a little mercury vapour in the vacuum of D. (3)
 - (c) (i) Draw a labelled diagram of a cathode ray oscilloscope (C.R.O) (3)
 - (ii) Describe how an alternating p.d may be measured using a C.R.O (3)

(d) In a Millikan's oil drop experiment a single negatively charged drop of radius 6×10^{-6} m was found to fall under gravity at a terminal velocity of 0.004 cm s⁻¹ and to rise at 0.012 cm s⁻¹ when a field of 2×10^5 Vm⁻¹ was suitably applied. Given that the viscosity of the medium was 2.122×10^{-5} Nsm⁻² determine the number of electrons on the drop. (5)

END

Name	Personal No
Signature	•••••

UGANDA ADVANCED CERTIFICATE OF EDUCATION MOCK ONE (1), 2016

PHYSICS PAPER 1

2 hours 30 minutes.

INSTRUCTIONS TO CANDIDATES:

Attempt **five** questions, including at least **one**, but not more than **two** from each of the sections **A**, **B**, and **C**.

Assume where necessary:


Acceleration due to gravity, g	=	9.81 ms ⁻²
Electron charge, e	=	1.6 x10 ⁻¹⁹ C
Stefan's Boltzmann's constant, σ	=	$5.7x10^{-8} \text{ Wm}^{-2} \text{ K}^{-4}$
Radius of the earth	=	6.4 x 10 ⁶ m
Radius of the sun	=	7.0 x 10 ⁸ m
Radius of Earth's orbit about the sun	=	1.5 x10 ¹¹ m
Speed of light in a vacuum, c	=	$3.0 \times 10^8 \text{ms}^{-1}$
Specific heat capacity of water	=	4,200J kg ⁻¹ K ⁻¹
Universal gravitational constant, G	=	$6.67 \text{ x} 10^{-11} \text{N m}^{-2} \text{ kg}^{-2}$
Avogadro's number N _A	=	$6.02 \times 10^{23} \text{ mol}^{-1}$
Density of water	=	1000 kgm ⁻³
Planck's constant, h	=	6.63 x 10 ⁻³⁴ Js
Charge to mass ratio, e/m	=	$1.8 \times 10^{11} \text{C kg}^{-1}$
The constant $^{1}/_{4\pi\varepsilon_{o}}$	=	9.0 x 10 ⁹ F ⁻¹ m

SECTION A

1. (a) Write the equations of *uniformly accelerated* motion (3 marks) (b) i) Briefly distinguish between *conservative* and *non-conservative* forces. Give two examples in each case. (4 marks) (ii) State the *principle of conservation* of energy. Show that a projectile fired with a speed $u ms^{-1}$ at an angle θ to the horizontal obeys this principle throughout its motion. (4 marks) (c) The engine of a 5-tonne lorry can develop 30kW, and its maximum speed on level road is 90 kmh^{-1} . Assuming that the friction resistance is constant, calculate the greatest speed at which the lorry can climb a hill of 1 in 25. (4 marks) (d) Define *moment* of a force. (i) (1 mark) A wheel of radius 0.84m is pivoted at its centre. A tangential force of 50 N (ii) acts on the wheel so that the wheel rotates with uniform velocity. Find the work done by the force to turn the wheel through 20 revolutions. (4 marks) 2. (a) (i) Define the term angular velocity. (1mark) (ii) A car of width c and whose centre of gravity is at a height h above the ground goes round a bend of radius r. Show that it will overturn if its speed exceeds $\sqrt{\frac{c g r}{2h}}$. (5 marks) (iii) A bucket of water is swung at constant tangential speed in a vertical circle of radius 0.5 m in such a way that the bucket is upside down when it is at the top of the circle. Find the minimum speed that the bucket may have if the water is to remain in it? (3marks) (b) State Kepler's laws of gravitation. (3 marks) (c) (i) Define gravitational field strength. (1mark)


- (ii) With aid of a sketch, qualitatively explain the variation of acceleration due to gravity with distance from the centre of the earth. (3marks)
- (d) A satellite of mass m is to be launched into an orbit of radius R round the earth of radius r. Show that the minimum launching velocity is given by $v^2 = \frac{g \, r (2 \, R r)}{R}$. (4 marks)
- 3 (a)(i) What is meant by the term **dimensions of a physical quantity**? (1 mark)
- (b) The volume per second of a liquid ejected from a pipe of radius a under steady flow is given by the expression $\frac{V}{t} = \frac{\pi}{8} p^x \eta^y a^z$, where p is the pressure gradient with dimensions $M L^2 T^{-2}$ and $[\eta] = M L^{-1} T^{-1}$. By dimensional analysis, find the values of x, y and z. (5 marks)
- (c) (i) Define the terms *range* and *time of flight* as applied to projectile motion. (2marks)
 - (ii) A shell is fired from a gun with a velocity U at an angle α to the horizontal. Show that the vertical distance y covered is given by $y = x \tan \alpha \frac{gx^2 \sec^2 \alpha}{2U^2}$. (4 marks)
- (d) A shell is fired with a velocity of $100 \, ms^{-1}$ at an angle of 60° to the horizontal from a gun placed at the top of a hill of vertical height 800 m above the ground. Find the:
- (i) time taken by the shell to hit the enemy on the ground assuming the shell is on target. (4 marks)
- (ii) velocity with which the shell hits the target. (4 marks)
- 4. (a)(i) Distinguish between **scalar** and **vector** physical quantities. Give **two** examples of each. (4marks)

(ii) The diagram below shows three coplanar forces acting on a particle.



Find the magnitude, and inclination to the 8N force, of the resultant force. (5marks)

- (b) State the laws of friction. (3marks)
- (c) With aid of a labeled diagram, describe an experiment to determine the coefficient of static friction. (4marks)
- (d) The figure below shows two particles M and N each of mass 2 kg. M is held at A on an inclined plane and is connected to N by a light inelastic string that passes over a smooth pulley P. The coefficient of friction between particle M and the plane is 0.5.



Find the acceleration of the masses and the tension in the string when M is released from rest. (4marks)

SECTION B

- 5. (a) (i) What is meant by a *thermometric property* and *triple point of water*? (2 marks)
 - (ii)Describe how you would measure the temperature of a body on the thermodynamic scale using a thermocouple. (3 marks)
 - (iii) The resistance of a platinum wire at the triple point of water is 5.25Ω . What will be the resistance at $100^{\circ}C$? (2 marks)
- (b) The resistance R_t of a platinum wire varies with the gas temperature t 0C according to the equation $R_t = R_0 \left(1 + \alpha t + \beta t^2\right)$, where R_0 is the resistance at 0 0C and α and β are constants. For a platinum resistance wire thermometer, $\alpha = 4.13 \times 10^{-3}$ and $\beta = -6.4 \times 10^{-7}$ respectively. Calculate the reading of the platinum thermometer when the gas thermometer reads 1000 0C .
- (c) Define the term **specific latent heat of vapourisation**. (1 mark)
- (d) An electrical heater rated at 500 W is immersed in a liquid of mass 2.0 kg contained in a large thermos flask of heat capacity 840 J kg⁻¹at 28°C. Electrical power is supplied to the heater for 10 minutes. If the specific heat capacity of the liquid is $2.5 \times 10^3~J~kg^{-1}~K^{-1}$, its specific latent heat of vapourization is $8.54 \times 10^3~J~kg^{-1}$ and its boiling point is $78~^{0}~C$, estimate the amount of liquid which boils off.

State any assumptions made in your calculation. (7 marks)

- 6(a) (i) State **Boyle's law**. (1 mark)
 - (ii) The pressure P of an ideal gas of density ρ and mean square speed $\overline{c^2}$ is given by $P = \frac{1}{3}\rho \ \overline{c^2}$. Use this expression to derive Boyle's law. (3 marks)
- (b) Air contains approximately of 20% oxygen and 80% nitrogen. The relative molecular masses of oxygen and nitrogen are 32 and 28 respectively. Calculate the ratio of the:
 - (i) mean-square speed of oxygen to that of nitrogen. (3 marks)
 - (ii) partial pressure of oxygen to that of nitrogen in air. (4 marks)

- (c) A fixed mass of a gas is contained in a cylinder of volume V_1 and pressure P_1 . The gas is allowed to expand to a volume V_2 and pressure P_2 . Assuming that the temperature remained constant, show that the work done by the gas is given by $W = -P_1V_1\ln\frac{P_2}{P_1}$. (3 marks)
- (d) A litre of hydrogen gas at a temperature of 27°C and a pressure 10⁵ Nm⁻² expands isothermally until its volume is doubled and is then adiabatically expanded until the new volume is again doubled. Given that $C_p = 29.2 \ J \ mol^{-1} \ K^{-1}$,
 - (i) Show the above processes on a P-V diagram, (2 marks)
 - (ii) Calculate the final pressure of the gas. (4 marks)
- 7(a) (i) Define **thermal conductivity** of a material (1 mark)
 - (ii)Briefly account for the fact that metals are better conductors of heat than insulators.

(3 marks)

- (b) A wall of a building consists of two brick layers each of thickness 10.0 cm and between which there is a layer of air 2.0 cm thick. Find the rate of heat flow through one m² of the wall if the inner and outer temperatures of the building are 25°C and 15°C respectively. (6 marks)
- (c) Describe with the aid of a labeled diagram, how the temperature of a furnace may be measured. (5 marks)
- (d) The total power output of the sun is $4.0\times10^{26}\,W$. Given that the mass of the sun is $1.97\times10^{30}\,kg$ and its density is $1.4\times10^3\,kg\,m^{-3}$, estimate the temperature of the sun. State any approximations made. (5 marks)

SECTION C

- 8(a) (i) Sketch the I-V characteristic for gaseous conduction. (2 marks)
 - (ii) Explain the main features of the characteristic. (3 marks)

(b)	(i) What is meant by thermionic emission?	(1 mark)
	(ii) Describe an experiment to show that cathode rays convey negative char	ge.
		(4 marks)
(c)	A horizontal beam of electrons, moving with a uniform speed, enters a uni- electric field between two horizontal parallel charged plates.	form
	Show that the path of the beam between the plates is a parabola.	(4 marks)
(d)	In a Millikan's oil drop experiment a single negatively charged drop of radiom was found to fall under gravity at a terminal velocity of $0.004~\rm cm~s^{-1}$ and $0.012~\rm cm~s^{-1}$ when a field of $2~\rm x~10^5~Vm^{-1}$ was suitably applied. Given that the of the medium was $2.122~\rm x~10^{-5}~Nsm^{-2}$ determine the number of electrons of	to rise at ne viscosity
9(a)	(i) Distinguish between X-rays and cathode rays.	(2 marks)
	(ii) In an X-ray tube explain the features adopted for the structure and mate the anode.	erials of (4 marks)
(b)	(i) State Bragg's law.	(1 marks)
	(ii) What is the condition for obtaining many orders of X-ray diffraction.	(1 marks)
	(iii) A monochromatic beam of X-rays of wavelength 1.187×10^{-10} m is incident set of parallel atomic planes of spacing 3.00×10^{-10} m. Determine the maximum of diffraction.	
(c)	(i) In photo-electricity, what is meant by the work function?	(1 mark)
	(ii) Describe how you could determine Planck's constant in a school laborate	tory. (5marks)
	(iii) When monochromatic light of frequency 6.0×10^{14} Hz falls on a metal stopping potential is 0.6 V while when the same surface is struck by light of frequency 1.0×10^{15} Hz the stopping potential becomes 2.2 V .	

Determine the work function of the metal.

(3 marks)

10(a)	(i) Distinguish between radioactivity and nuclear fission .	(2 marks)
	(ii) Define binding energy of a nucleus	(1 mark)
(b)	(i) What is half life of a radioactive substance?	(1 mark)
	(ii) Derive the relationship between half life and the decay constant of a rasubstance.	ndioactive (3 marks)
(c)	With the aid of a diagram describe how a Geiger-Muller tube detects ioni particles.	zing (6 marks)
(d)	At a certain time, an α-detector registered a count rate of 104 min ⁻¹ . Exac	tly 12 days
	later, the count rate dropped to 13 min ⁻¹ . Find the decay constant.	(4 marks)
(e)	Explain how each of the following would affect the rate of decay in (d)	
	(i) temperature rise	(1 mark)
	(ii) mass of the radioactive substance	(1 mark)
	(iii) exposed surface area of the radioactive substance	(1mark)

END

Name	Center/Index no
Signature	House

UGANDA ADVANCED CERTIFICATE OF EDUCATION

BEGINNING OF TERM 3, 2014

PHYSICS PAPER 1

2 hours 30 minutes.

9.81 ms⁻²

INSTRUCTIONS TO CANDIDATES:

Attempt five questions, including at least one, but not more than two from each of the sections A, B, and C.

Assume where necessary:

Acceleration due to gravity, g

Electron charge, e = 1.6 x10⁻¹⁹ C Stefan's Boltzmann's constant, σ = 5.7x10⁻⁸ Wm⁻² K⁻⁴ Specific heat capacity of copper = 400 J kg⁻¹ K⁻¹ Specific latent heat of fusion of ice = 3.36 x 10⁵ J kg⁻¹ Radius of Earth's orbit about the sun = 1.5 x10¹¹m Speed of light in a vacuum, c = 3.0 x 10⁸ms⁻¹

Specific heat capacity of water = $4,200 \text{J kg}^{-1} \text{ K}^{-1}$ Universal gravitational constant, G = $6.67 \times 10^{-11} \text{N m}^2 \text{ kg}^{-2}$ Avogadro's number N_A = $6.02 \times 10^{23} \text{ mol}^{-1}$

Density of water = 1000 kgm^{-3} Planck's constant, h = $6.63 \times 10^{-34} \text{ Js}$ Charge to mass ratio, e/m = $1.8 \times 10^{11} \text{ C kg}^{-1}$ The constant $1/4\pi\epsilon_0$ = $9.0 \times 10^9 \text{ F}^{-1}\text{m}$

Permittivity of free space, ϵ_0 = 8.85 x 10^{-12} Fm⁻¹ Mass of the electron = 9.11 x 10^{-31} kg

SECTION A

1(a) (i) Define a **couple.** (1)

(ii) A pair of forces of magnitude \mathbf{F} , act on a wheel and rotates it through an angle $\mathbf{\theta}$, show that the work done by the forces is equal to the product of torque and the angle of rotation? (3)




Fig. 1

The forces of **3.5N**, **2.5N** and **1.7N** act at a point as shown in figure 1. Find the resultant force. (6)

- (c) A motorist traveling at a constant speed of **50 kmhr** ⁻¹ passes a motorcyclist just starting off in the same direction. If the motor cyclist maintains a constant acceleration of **2.8 ms** ⁻², calculate;
 - (i) the time taken by the motor cyclist to catch up with the motorist. (4)

	(ii) the speed with which the motor cyclist overtakes the motorist.	(2)	
(d)	Explain why the tension in a cable of a lift is different when the lift is		
	ascending from when it is descending.	(4)	
2(a)	State Newton's law of motion	(3)	
(b)	Use Newton's laws of motion to show that linear momentum is conserved		
	when two parties collide directly.	(5)	ı
(c)	Three balls A, B and C of masses M_1 , M_2 , and M_3 respectively lie in a straight smooth surface. The balls are initially at rest. Ball A is projected with a veloci B, makes an elastic collision with B. If B moves and makes a perfectly inelastic C, Show that both B and C move with a velocity	ty, ${\it V}_1$ towards	
	$V_3 = \frac{2 M_1 M_2 V_1}{(M_1 + M_2) (M_2 + M_3)}$	(5)	
(d)	A bullet of mass 10g traveling horizontally at 100 ms ⁻¹ embeds itself		
	in the centre of a block of wood of mass 900g which is suspended by a		
	light vertical string 1m in length.		
	(i) Determine the vertical height through which the combination rises.	(4)	
	(ii) Calculate the loss in kinetic energy of the bullet.	(3)	
3(a)	(i) Define pressure.	(1)	

	(ii) Derive the expression for the pressure at a point a depth ${\bf h}$ from the		
	surface of a liquid of density ρ .	(3)	
(b)	A block of mass 0.10 kg is suspended from a spring balance. When the block immersed in water of density $1.0 \times 10^{3} \text{ kgm}^{-3}$, the spring balance reads the block is immersed in a liquid of unknown density, the spring balance Find;		When).70N.
	(i) the density of the solid.		(4)
	(ii) the density of the liquid.	(3)	
(c)	What is meant by the following terms as applied to fluids?		
	(i) terminal velocity.		(1)
	(ii) coefficient of viscosity.		(1)
	(iii) streamline flow.		(1)
(d)	Explain qualitatively the origin of the lift force on an aero -plane at take-off.	(3)	
(e)	Briefly explain, using the kinetic theory of matter, the effect of temperature of viscosities of fluids.	on	(3)

- 4(a) (i) Define *simple harmonic motion.* (1)
 - (ii) Give *four* characteristics of simple harmonic motion. (2)
- (iii) Give two examples of oscillatory motion which approximates to simple harmonic motion. (1)
- (b) Two springs of force constants K₁ and K₂ are suspended from a horizontal support. A mass m, hangs from the mower ends of the springs as shown in Figure 2.




Fig. 2

If both springs have negligible mass, show that when m is displaced from its equilibrium position, it describes simple harmonic motion. Hence deduce the expression for the frequency of the oscillations. (6)

(c) The figure 3 shows two springs S₁ and S₂ in series fixed between a wall and a particle of mass 0.3 kg which is resting on a smooth horizontal plane.




Fig. 3

	The force constants of S ₁ and S ₂ are 20 Nm ⁻¹ and 30 Nm ⁻¹ respectively. The horizontally through a distance of 4 cm from its rest position and then rele	_	∍d
	(i) the maximum tension in the springs.	(3)	
	(ii) the frequency of oscillation.	(2)	
	(iii) the displacement of the particle from the equilibrium position $0.4\mathrm{s}$ after (4)	release. 4)	
	SECTION B		
5(a)	(i) What is a thermometric property ?	(1)	
	(ii) Give two examples of thermometric properties	(1)	
	(iii) Describe how the platinum resistance thermometer can be used to me absolute temperature.	asure (6)	
(b)	(i) Define specific heat capacity.		(1)
	(ii) Cold water at 10℃ and hot water at 85℃ are run into a bath tub at a ra V m³ per minute and 3 ×10 -2 m³ per minute respectively. At the point of filling the tub, the temperature of the mixture was 35℃. Calculate the time taken to fill the tank if its capacity is 1.5m³	te of	(6)
(c)	State three advantages of the continuous flow method over the method of	ı	

	mixtures in the determination of specific heat capacity of a liquid.	(3)	
(d)	Explain why water is preferred for use in the cooling systems of engines to other liquids.		(2)
6(a)	(i) Define specific latent heat of vaporization.	(1)	
	(ii) State the factors that determine the rate of cooling of a body.	(2)	
	(iii) Explain why a small body cools faster than a bigger one of the same material and shape.		(4)
(b)	With the aid of a well labeled diagram, describe a method of determining the specific latent heat of vaporisation of water based on mixtures.	(7)	
(c)	Two taps, one delivering water at a temperature of 25 °C at a rate of 0.3gs ⁻¹ and the other delivering water at a temperature of 15 °C at a rate of 0.4gs ⁻¹ , are directed to a well lagged copper calorimeter of mass 50g containing ice at 0 °C while stirring. After 5 minutes the contents of the calorimeter are found to have attained a temperature of 5 °C. Find the		
	mass of ice originally in the calorimeter.		(6)

7(a)	(i) What is meant by an ideal gas?		(1)
	(ii) State three differences between a real and an ideal gas.	(3)	
	(iii) What is meant by kinetic theory of matter?		(2)
of mat	(iv) Describe briefly an experiment which one can carry out in support eter. (4)	of the kinetic	theory
(b) square	Derive the expression $P=\frac{1}{3}\rhoc^{\overline{2}}$ for the pressure of an ideal gas of c e-speed, $c^{\overline{2}}$.	lensity, ρ and (5)	mean-
	An ideal gas of volume 100 cm³ at s.t.p expands adiabatically until its properties original value. Find the new volume and temperature if the ic heat capacities is 1.4. (5)	ressure ratio of the pri	drops ncipal
	SECTION C		
8(a)	(i) State Rutherford's model of the atom.		(2)
	(ii) Explain two main failures of Rutherford's model of the atom.		(3)
(b)	(i) Explain how Millikan's experiment for measuring the charge of the charge is quantized.	electron prov	es that

(c)	In the measurement of electron charge by Millikan's apparatus, a potential difference is switched on, the drop rises with constant velocity of 4.0× 10 ⁻⁴ to the potential difference is switched on, the drop rises with constant velocity of 1. If the mass of the oil drop is 1.0× 10 ⁻¹⁴ kg, find the number of electron of drop.(Assume air resistance is proportional to the velocity of the oil drop and rethrust due to air)	ifference ns ⁻¹ . When 8.0 × 10 ⁻⁵ ms ⁻ charges on the
(d)	Explain qualitatively, the use of the time-base in a cathode –ray oscilloscope.	(4)
9(a)	(i) Distinguish between radioactivity and nuclear fission.	(2)
(2)	(ii) Why are neutrons preferred to charged particles for inducing nuclear	reactions.
(b)	Describe how a Geiger-Muller tube works in the detection of ionizing radiatio	n. (6)
(c)	1.2g of a substance form a point source of γ -rays. Only 1 in 10^{12} of its atoms are radioactive and the half-life is 100 days. A Geiger-Muller tube facing the source at a distance of 10 cm gives a count rate of 11 s ⁻¹ . If the window of the tube has an area of 7 cm^2 , find;	
	(i) the number of radioactive atoms present.	(5)
	(ii) the mass number of the substance.	(2)
(d)	Determine whether the nucleus $^{210}_{84}$ <i>Po</i> is stable or it may undergo	

disintegration to produce $^{206}_{82}$ *Pb* and an α -particle. (3)

Mass of ${}^{210}Po = 209.937u$

Mass of $^{206}Pb = 205.929u$

Mass of ${}^{4}\text{He} = 4.002u$

- 10(a) (i) What are Positive rays? (1)
 - (ii) State four properties of Positive rays. (2)
- (b) With the aid of a diagram, describe the operation of the Bainbridge spectrometer in determining the specific charge of ions. (5)
- (c) Show that when an alpha particle of mass, m, and initial velocity, V, collides head-on with an atom of atomic number, Z, the closest distance of approach to the nucleus is

$$a_0 = \frac{Z e^2}{\pi \epsilon_0 m V^2}$$

Where, e is the electronic charge and ϵ_{o} is the permittivity of free space.

(4)

- (d) (i) Derive Bragg's law of X-ray diffraction. (4)
- (ii) Calculate the atomic spacing of sodium chloride if the relative atomic mass of sodium is 23.0 and that of chlorine is 35.5. [Density of sodium chloride is 2180 kgm⁻³] (4)

END

Name	Center/Index no
C: 4	
Signature	

UGANDA ADVANCED CERTIFICATE OF EDUCATION MOCK ONE (1), 2014

PHYSICS PAPER 1

2 hours 30 minutes.

INSTRUCTIONS TO CANDIDATES:

Attempt **five** questions, including at least **one**, but not more than **two** from each of the sections **A**, **B**, and **C**.

Assume where necessary:

Acceleration due to gravity, g = 9.81 ms^{-2}

Electron charge, $e = 1.6 \times 10^{-19} \text{ C}$

Stefan's Boltzmann's constant, $\sigma = 5.7x10^{-8} Wm^{-2} K^{-4}$

Radius of the earth = $6.4 \times 10^6 \text{m}$

Radius of the sun $7.0 \times 10^8 \text{m}$ Radius of Earth's orbit about the sun $1.5 \text{ x} 10^{11} \text{m}$ Speed of light in a vacuum, *c* $3.0 \times 10^8 ms^{-1}$ Specific heat capacity of water 4,200J kg⁻¹ K⁻¹ Universal gravitational constant, G $6.67 \ x10^{-11} N \ m^{-2} \ kg^{-2}$ Avogadro's number NA 6.02 x 10²³ mol⁻¹ 1000 kgm⁻³ Density of water $6.63 \times 10^{-34} \text{ Js}$ Planck's constant, h 1.8 x 10¹¹ C kg⁻¹ Charge to mass ratio, e/m $^{1}/_{4\pi\varepsilon_{o}}$


SECTION A

9.0 x 10⁹ F⁻¹m

1. (a) (i) State the laws of solid friction. (3)

The constant

- (ii) Explain what is meant by the angle of friction. (2)
- (b) Describe an experiment you would perform to determine the coefficient of static friction between a sphere and a given plane surface. (4)
- (c) Figure 1 shows a body in form of a uniform cube of weight 200 N resting with one of its edges B on a horizontal floor and edge A against a vertical wall.



B **Fig. 1**

When a boy of weight 100 N sits on edge D, the cube rests in limiting equilibrium with face AB making an angle of 30° with the horizontal. Given that the coefficient of static friction is the same at every point of contact find

(i) the coefficient of static friction. (7)

(ii) the reactions at the wall and the floor. (4)

2. (a) (i) Define *coefficient of viscosity* of a liquid. (1)

(ii) A ball of density σ and radius r is released to fall through a liquid viscosity, η . Derive an expression for its terminal velocity during its (3)	•
(iii) Describe a simple experiment to demonstrate stream line and turn a liquid. (4)	bulent flow in
(b) (i) State Bernoulli's principle.	(2)
(ii) Explain one application of Bernoulli's principle.	(3)
(c) (i) Given that the surface tension of a soap solution is 2.7 x 10 ⁻² Nm work done in forming 10 soap bubbles each of diameter 1.6 cm. (3)	1 ⁻¹ , calculate the
(ii) Two soap bubbles, one of radius x and the other of larger radius y attached to each other. Find an expression for the radius of the com-	_
interface in terms of <i>x</i> and y.	(4)
3. (a) (i) State Newton's law of gravitation.	(1)
(ii) Deduce the dimensions of the Universal gravitational constant.	. (2)
(b) Assume that a planet moves with uniform speed in a circular orbit the sun. The periodic time of the planet is T. Show that the quantity independent of the mass of the planet. (4)	

(c) An artificial satellite is launched in a circular orbit round the equator at a of 3.60 x 10^7 m above the surface of the Earth.	height	
(i) Determine, from first principles, the speed with which the satellite mulaunched to maintain it in the orbit. (5)	st be	
(ii) Determine the periodic time of the satellite. (3))	
(iii) What deduction can be made from the result obtained in (ii) above wit reference to the satellite? (2)	:h	
(iv) If another satellite moves in a circular orbit concentric with that of the satellite, at a height of 1.48×10^{-7} m above the surface of the Earth, what is the time of this second satellite? (3)		
4. (a) Write the equations of uniformly accelerated motion. (3)	
(b) A reckless driver traveling at $40~\rm ms^{-1}$ and accelerating at $0.2~\rm ms^{-2}$ overtakes a police car at the instant when the police car starts up in pursuit. If the police car maintains a uniform acceleration of $1~\rm ms^{-2}$, calculate;		
(i) how long the police car takes to catch up with the car. (3))	
(ii) the distance the police car travels before catching up with the car. (2))	

(c) (i) Briefly distinguish between conservative and non-conservative forces an example of each. (4)	s. Give
(ii) State the principle of conservation of energy. Show that a projectile fired upwards obeys this principle throughout its downward motion. (4)	l
(d) The engine of a 5-tonne lorry can develop 30kW, and its maximum speed road is 90 kmh ⁻¹ . Assuming that the friction resistance is constant, calculat speed at which the lorry can climb a hill of 1 in 25 (4)	on level te the greatest
SECTION B	
5. (a) With the aid of a diagram, describe how you would detect infra-red beam of sunlight. (4)	radiation in a
(b) (i) Sketch curves to show how energy is distributed among the various of black body radiation for three different temperatures.	wavelengths
(ii) Explain why cavities in a fire look brighter than the rest of the fire.	(2)
(iii) The intensity of radiant energy from a black body is a maximum at a wavelength of 1.5 x 10 ⁻⁶ m. Calculate the temperature of the black body	7. (2)
(c) (i) State Stefan's law of black body radiation.	(1)
(ii) A strip of platinum foil coated black, is placed on the ground. The are	ea of the

strip exposed to the radiation from the sun is 10^{-2} m². The radiation

from the sun falls

normally on the strip to obtain a certain temperature the radiation and a current 1.4 A is maintained through t 4.0 V to obtain the same temperature rise. If only 40% incident on the earth's atmosphere reaches the earth's temperature of the sun.	the foil at a potential difference of of the intensity of radiation
	• •
(iii) If the foil in (ii) were not painted black but was be applied across it to maintain a constant temperature?	80% black, what p.d would have to (2)
6. (a) (i) Define the specific latent heat of vaporisation of	a substance. (1)
(ii) Describe an experiment to determine the specific by the electrical method. (5)	latent heat of vaporisation of water
(b) (i) Explain why the molar heat capacity of an ideal from that at constant volume.	gas at constant pressure differs (3)
(ii) Derive an expression for the difference of the abo	ove molar heat capacities. (4)
(c) (i) State the conditions necessary for a reversible ad	iabatic process. (2)
(ii) A gas initially occupying a volume of 1.0 litre at	a temperature of 47°C and a
pressure of 1.40 x 10 ⁵ Pa expands isothermally to a volu	
compressed adiabatically to its original volume.	If γ = 1.4, find the final
temperature and pressure. (5)	

7. (a) What assumptions are necessary in the derivation of the Kinet	ic theory
expression for the pressure of an Ideal gas?	(4)
(b) A beam of 2 x 10^{22} Nitrogen atoms, each of mass 2.32 x 10^{-26} normally on a wall of a cubical container of edge 5.0cm. The beam is 180°. If the mean speed of the atoms is 500ms ⁻¹ , find the pressure exert gas. (4)	reflected through
(c) (i) State Dalton's law of partial pressures.	(1)
(ii)Two containers A and B of volume 3 x 10 ³ cm ³ and 6 x 10 ³ cm ³ rescontain helium gas at a pressure of 1.0x10 ³ Pa and temperature 300 is heated to 373K while container B is cooled to 273K. Find the final helium gas. (5)	
(d) Use the kinetic theory of gases to explain the effect on a satura pressure of;	ted vapour
(i) raising the temperature at constant volume.	(3)
(ii) increasing the volume at constant temperature.	(3)
SECTION C	
8. (a) Draw a set-up of the apparatus used in the famous Millikan's oil and answer the following questions: (2)	drop experiment

(i) \	Why should a constant temperature bath be necessary?	$^{\prime}$ (2)
-------	--	-----------------

(b) The diagram shows an ultraviolet lamp L placed near a clean but insulated zinc plate P in a dark room. With L switched off, the plate is negatively charged and connected to the cap of an electroscope as shown.





Fig. 2

Explain what is observed when;

(c) An X-ray tube works on a d.c potential of 50 kV. Only 0.4% of t	the energy	of the
cathode rays is converted into X-radiation and heat is generated in	the target at a	rate of
600 W. Estimate;		

- (i) the number of electrons striking the target per second. (3)
- (ii) the velocity at which the electrons strike the target. (2)
- 9. (a) Explain the origin of the **continuous** and **characteristic line** spectra in an X-ray tube. (5)
- (b) A monochromatic beam of X-radiation is incident on a set of parallel atomic planes of sodium chloride (NaCl) crystal. A first order diffraction maximum is observed at a glancing angle of 5.4°. When the same X- radiation is incident on a similar set of atomic planes of a potassium chloride (KCl) crystal the second order diffraction maximum is observed at a glancing angle of 10.2°. Compare the spacing between the respective atomic planes for the two crystals.
 - (c) The diagram below depicts possible electron orbits in the Bohr model for hydrogen. Assume the orbits are circular





n = 1

Fig. 3

(i) Show that the total energy of an electron in an orbit of radius r is given by $E = -e^2$.

 $8\pi\epsilon r$ (4)

(ii) If only the orbits allowed for are those whereby

$$mvr = \underline{nh},$$
 2π

where m is the electronic mass, v the electron speed, n an integer and h Planck's constant, show that the total energy in (i) above can be expressed as $E_n = -me^4$.

$$8\varepsilon_o^2 h^2 n^2 \tag{3}$$

- (iii) Calculate the wavelength of the radiation that will be emitted when the electron makes a transition from n = 4 to n = 2 orbits. (5)
- 10. (a) (i) What is meant by the term 'binding energy of a nucleus'? (1)
- (ii) Calculate the binding energy per nucleon of an α -particle, expressing your result in **MeV**.

Mass of proton	= 1.0080u	
Mass of a neutron	= 1.0087u	
Mass of an α - particle	= 4.0026u	
1u	= 931 MeV. ((4)

(iii) Sketch a graph of binding energy per nucleon against mass number and use it to explain liberation of energy by nuclear fusion and nuclear fission.

(5)

- (b) Derive an expression relating the half life of a radioactive material, $T_{\frac{1}{2}}$ and the decay constant, λ .
- (c) When $^{238}_{92}$ U decays, the end product is $^{206}_{82}$ Pb. The half life is 1.4×10^{17} s.

Suppose a rock sample contains $^{206}_{82}Pb$ and $^{238}_{92}U$ in the ratio 1: 5 by weight. Calculate the:

- (i) number of $^{206}_{82}$ Pb atoms in 5.0 g of the rock sample. (3)
- (ii) age of the rock. (4)

END

Name	Center/Index no
Signature	House

UGANDA ADVANCED CERTIFICATE OF EDUCATION

MOCK TWO (2), 2014

PHYSICS PAPER 1

2 hours 30 minutes.

INSTRUCTIONS TO CANDIDATES:

Attempt five questions, including at least one, but not more than two from each of the sections A, B, and C.

Assume where necessary:

Acceleration due to gravity, g = 9.81 ms^{-2} Electron charge, e = $1.6 \times 10^{-19} \text{ C}$

Stefan's Boltzmann's constant, σ = 5.7x10⁻⁸ Wm⁻² K⁻⁴

Radius of the earth $= 6.4 \times 10^6 \text{m}$ Radius of the sun $= 7.0 \times 10^8 \text{m}$ Radius of Earth's orbit about the sun $= 1.5 \times 10^{11} \text{m}$ Speed of light in a vacuum, $c = 3.0 \times 10^8 \text{ms}^{-1}$ Specific heat capacity of water $= 4,200 \text{J kg}^{-1} \text{ K}^{-1}$

Universal gravitational constant, G = $6.67 \times 10^{-11} \text{N m}^2 \text{ kg}^{-2}$ Avogadro's number N_A = $6.02 \times 10^{23} \text{ mol}^{-1}$

Density of water = 1000 kgm^{-3} Planck's constant, h = $6.63 \times 10^{-34} \text{ Js}$ Charge to mass ratio, e/m = $1.8 \times 10^{11} \text{ C kg}^{-1}$ The constant $1/4\pi\epsilon_0$ = $9.0 \times 10^9 \text{ F}^{-1}\text{m}$

Permittivity of free space, ϵ_0 = 8.85 x 10^{-12} Fm⁻¹ Mass of the electron = 9.11 x 10^{-31} kg

	SECTION A		
1(a)	What is meant by the following terms as applied to materials?		
	(i) tensile stress	(1)	
	(ii) tensile strain	(1)	
	(iii) elastic limit	(1)	
(b) (3)	Explain the occurrence of work hardening in an otherwise ductile	mate	rial.
(c)	Describe an experiment to determine Young's modulus for a steel win	re. (5)	
	A rod of cross section area A and Young's modulus E is fixed in rigid Given that the linear expansivity of the rod is α derive an expression in it if the temperature changes by $\delta\theta$. (3)		orts at its e force
point	A metal wire of diameter 2.0 x 10 ⁻⁴ m and length 2 m has its ends fill B 1.6 m apart, AB being horizontal. What force should be applied of the wire to pull the point of application to a point 0.65 m below $lus = 2 \times 10^{11} \text{ Nm}^{-2}$ (6)	ed at th	
2 (a)	(i) What are the distinguishing characteristics between <i>uniform circumotion</i> and <i>simple harmonic motion</i> ?	lar (3)	
radius	(ii) Derive an expression for the acceleration of a particle describing a s r with a uniform speed, v. (5)	circle	of
	A particle of mass m is hung on a string of length, l. It is then projected ontal velocity, V.	ed	with a
subse	(i) Show that when the string makes an angle θ with the vertical, during quent motion, the tension in the string is	ing	the
	$m\{V^2/I - g(2 - 3cos\theta)\}$	(4)	
	(ii) Hence show that if the particle is to describe a complete circle, the	en	
	$V^2 > 5gI$	(3)	
(c)	The displacement x of a particle is given by $x = a \sin \omega t$		
	(i) Derive an expression for its velocity in terms of its displacement,		
	angular velocity, ω , and the amplitude, a .	(3)	
	(ii) Show that the total energy of the particle is constant.	(2)	

3(a)	(i) What is meant by acceleration due to gravity?	(1)	
(1)	(ii) Sketch the displacement-time graph for a body thrown vertically up	p in	air.
	(iii) What is meant by a trajectory?	(1)	
air at a	(iv) Show, from first principles, that the path followed by a particle angle, α , to the horizontal is parabolic. (3)	projec	ted in
(b)	A projectile is fired horizontally from the top of a cliff, 200 m high. The projectile lands 1000 m from the bottom of the cliff. Find the:	he	
	(i) Initial speed of the projectile.	(4)	
	(ii) Velocity of the projectile just before it hits the ground.	(5)	
(c) helica	Describe an experiment to determine the acceleration due to gravity ull spring of unknown force constant. (5)	sing	a
4(a)	(i) Account for the existence of intermolecular forces.	(2)	
substa	(ii)Sketch a graph of potential energy against separation of two molecunce and explain the main features of the graph. (4)	les	in a
(b)	(i) Define surface tension.	(1)	
	(ii)Show that the excess pressure, p, in an air bubble inside a liquid ove outside pressure is given by	r	

 $p = 2\gamma$

r

	where $ r $ is the radius of the bubble and $ \gamma $ its surface tension.	(3)
(c)	(i) State Archimedes principle.	(1)
of the	(ii) Show that the Up thrust on a body immersed in a fluid is equal to the fluid displaced. (4)	e weight
(d)	A solid weighs 237.5g in air and 12.5g when totally immersed in a liquid relative density 0.9. Calculate;	l of
	(i) the relative density of the solid	(2)
its volu	(ii) the relative density of a liquid in which the solid would float with or time exposed above the liquid surface. (3)	ne- fifth of
	SECTION B	
5(a)	Define <i>thermo- conductivity</i> of a material.	(1)
(b)	(i) Explain, using the molecular theory of matter the mechanism of heaconduction in insulators.	at (3)

insula	(ii) Briefly account for the fact that metals are better conductors tors. (3)	s of heat	than
(c) betwe	A wall of a building consists of two brick layers each of thicknessen which there is a layer of air 2.0 cm thick.	ess 10.0 cm	and
	Find the rate of heat flow through 1 m ² of the wall if the inner a temperatures of the building are 25°C and 15°C respectively.	and outer	
	[Thermal conductivities of brick and air are respectively, 0.7 W	m-1K-1 and	
	$0.024~\mathrm{Wm^{-1}K^{-1}}]$	(6)	
(d)	(i) What is meant by a black body?	(1)	
	(ii) The average distance of Pluto from the sun is 40 times that on. If the sun radiates as a black body at 6000K, and is calculate the surface temperature of Pluto. (6)	of the earth 1.5×10^{11} m f	from from the
6(a)	(i) What is meant by cooling correction?	(1)	
heat c	(ii) Explain how the cooling correction may be estimated in the apacity of poor conductor of heat by the method of mixtures.		n of the
(b)	(i) Define the term specific heat capacity of a substance.	(1)	

power	(ii)An electrical heater rated 500W is immersed in a liquid of mass 2.0 k contained in a large thermos flask of heat capacity 840J K ⁻¹ at 28 ⁰ C. It is supplied to the heater for 10 minutes. If the specific heat capacity of	Electrical f the liquid is
	0 ³ J kg ⁻¹ K ⁻¹ , its specific latent heat of vaporization is 8.54 x 10 ³ J kg	_
	g point is 78°C, estimate the amount of liquid which boils off stating any	assumptions
made.	(7)	
(c)	(i) With reference to an electrical thermometer, describe the steps involved	ved in
setting	up a Kelvin scale of temperature. (3)	
	(ii)The resistance of the element of a platinum resistance thermometer is	
	1	on the
platinu	Im resistance scale would correspond to a resistance of 9.84Ω ?	
	(3)	
7(a)	Explain why	
	(i) at low pressures a real gas approximates to a perfect gas.	(2)
		· /
	(ii) evaporation causes cooling.	(2)
(b)	(i) State three properties of a saturated vapour.	(3)
	(ii) Air is trapped between a piston and a liquid in a cylinder. The piston	·
=	-	the liquid
	aporated. Sketch a graph of pressure against volume for the conten	ts of the
cylinde	er and explain its features. (3)	

(iii) Describe an experiment to show that saturated vapour pressure va temperature. (6)	ries with
(c) The saturated vapour pressure of ether at 0° C is 185 mmHg and at 20 440 mmHg. The bulb of a constant-volume gas thermometer contains dry at ether for saturation. If the observed pressure in the bulb is 1000 mmHg will it be at 0° C? (4)	ir and sufficient
SECTION C	
8(a) Define the following terms as used in photo electricity.	
(i) work function.	(1)
(ii) stopping potential	(1)
(b) (i)Describe an experiment to determine Planck's constant in a laborate	tory. (5)
(ii) Write down Einstein's photo-electric equation and explain how it emission of electrons from a metal surface. (3)	accounts for

	magnetic radiation of frequency .5eV. Calculate the maximum s	=	
from the surface.	(4	·	
(c)	P	_	
		_ Q	
		4	Fig.1
			118.1
separated by a distribution between P and Q i directed midway b	nows two parallel metal plates P ance of 4.0 cm. A p.d. of 12V is s a vacuum. A beam of electron etween P and Q. Calculate P and Q to the initial dire	applied between P s of speed u = 1.0 the angle at which th	and Q and the space 0 x 10 ⁶ ms ⁻¹ is
9(a) (i) Distingu	aish between X-rays and cathod	e rays.	(2)
(ii) Draw a production of the 2	well-labeled diagram of the X-ra X-rays.	ay tube and outline th	ne principles of
(b) (i) State Br	agg's law of X-ray diffraction.		(1)

set of 1	(iii) A monochromatic beam of X-ra parallel atomic planes of spacing 3.00	•	-	
diffrac				
(c)	Briefly distinguish between X-ray l	line spectra a	nd optical line spectra. (2)
(d)	Figure 2 below shows some of the e	energy levels	of a mercury atom	
	Energy in eV			
	0.0	Ε _σ	2	
	-1.6			
	-3.7	E	3	
	-5.5	Е		
	-10.4	Eı	ı	
		Fig. 2		
	Determine			
	(i) the minimum speed of an electro	n that could	just ionize the mercury	atom.
(3)				
E2.	(ii) the wavelength of the radiation	emitted in a	n electron transition from (3)	n E ₃ to
Li2.			(0)	

- 10(a) (i) Sketch and label the I-V characteristic for gaseous conduction. (2)
 - (ii) Explain the main features of the curve. (3)
- (b) A valve diode D was connected to rectify a.c supply to a load as shown in figure 3. C is a capacitor.




Fig. 3

- (i) State the purpose served by the capacitor C. (1)
- $\label{eq:continuous} \mbox{(ii) Explain the effect on the p.d's V_d and V_L of introducing a little } \qquad \mbox{mercury} \\ \mbox{vapour in the vacuum of D.} \qquad \qquad \mbox{(3)}$


(c)	A sinusoidal voltage of amplitude 0.1V is applied to the grid of triode of amplification factor 12. If the anode resistance of the triode is 15 k Ω , Find;		
	(i) Voltage will appear across a load of 18 $\mbox{k}\Omega$	(3)	
	(ii) The gain of the stage at this load resistance.	(2)	
(d)	(i) Draw a labeled diagram of a cathode ray oscilloscope (C.R.O)	(3)	
	(ii) Describe how an alternating p.d may be measured using a C.R.O	(3)	
	END		
	S.6 PHYSICS PAPER 1 – TEST 2		
1	(a) (i) Define surface tension and show that it is equal to the work dor in increasing the liquid surface under isothermal conditions.	ne per unit area (3)	
	(ii) Find the work required to break up a drop of water of radius 0.5 cr water each of radius 1mm . [Surface tension of water is 2.27 x 10 ⁻² N	-	
	(b) Describe an experiment to measure surface tension of a liquid by th tube method.	e capillary (6)	

- (c) (i) Derive an expression for the excess pressure in a soap bubble of radius r if the soap solution has surface tension γ . (3)
- (ii) The pressure inside an air bubble formed 40 cm below the water surface is

1.063 \times **10**⁵ **Pa**. If the surface tension of water is **2.27** \times **10**⁻² **Nm**⁻¹ find the radius of the bubble. [Atmospheric pressure = **1.013** \times **10**⁵ **Pa**] (4)

2. (a) State the characteristics of photoelectric emission. (4)

(b) The figure below shows graphs of current I against p.d V applied across a photoelectric cell that receives monochromatic light for two different intensities A and B (A < B)



-V V_T +V

- (i) Draw a diagram to show the set-up you could use to establish these results. (3)
- (ii) State what V_T represents and explain the features of the graphs. (3)

- (c) Describe an experiment to determine Planck's constant in a laboratory. (5)
- (d) Light of frequency **5.0** \times **10**¹⁴ **Hz** liberates electrons with maximum energy **2.31** \times **10**⁻¹⁹ **J** from a certain metallic surface. What is the wavelength of ultra-violet light which liberates electrons of maximum energy **8.93** \times **10**⁻¹⁹ **J** from the surface? (5)

FND

(ii) A cylinder contains 100 litres of gaseous oxygen at a pressure of 1.217×10^7 Pa and temperature 20° C. Assuming oxygen behaves as an ideal gas in this region of pressure and temperature, find the volume of liquid oxygen (density 1140 kg m⁻³) that may be made by liquefying completely the contents of the cylinder.

[Relative molecular mass of oxygen = 32]

(4)

(c) In a continuous flow method for determining the specific heat capacity of a liquid the following data were obtained.

P.d/V	Current/A	Mass of liquid collected in 15 min /kg	Inflow temperature /°C	Outflow temperature /°C
6.0	2.5	0.220	25.0	35.0
7.2	3.0	0.366	25.0	35.0

(i) Draw a labelica diagram of the apparatas necaea to obtain sach data.	(i)	Draw a labelled diagram	of the apparatus needed to obtain such data.	(4
--	-----	-------------------------	--	----


(ii) Calculate the specific heat capacity of the liquid.

(3)

(iii) Find the rate at which heat is lost to the surroundings.

(2)

- (c) Two enemy artillery guns P and Q are 4 km apart. P fires a shell at 200 ms⁻¹ while Q fires one at 240 ms⁻¹, each to destroy the enemy gun. If the shells are fired simultaneously and each gun is successfully hit, find
 - (i) the angle of projection of each shell (3)
 - (ii) when one is hit, the time that elapses before the other is hit (3)
- (iii) the horizontal distance moved by each shell by the time one shell is vertically above the other. (4)
- (b) A sphere of mass 2 kg moving with a velocity of 3 ms⁻¹ impinges directly on another of mass 1 kg moving at 2 ms⁻¹ in the opposite direction. If the coefficient of restitution is 0.5, determine the velocity of each sphere just after impact. (5)
- (c) The figure shows two particles M and N each of mass 2 kg. M is held at A on an inclined plane and is connected to N by a light inelastic string that passes over a smooth pulley P.



The coefficient of friction between particle M and the plane is 0.5.

- (i) If M is released, find the acceleration of the masses and the tension in the string (6)
- (ii) If, during motion, the inclination of the plane is varied, at what inclination to the horizontal will the system begin to move with constant speed? (3)
- (b) A mass of 0.1kg suspended from a spring of force constant 24.5 Nm⁻¹ is pulled vertically downwards through a distance of 5.0cm and released. Find the

- (i) period of oscillation (2 marks)
- (ii) position of the mass 0.3 seconds after release (4 marks)

Name	Personal no
Signature	House

UGANDA ADVANCED CERTIFICATE OF EDUCATION

BEGINNING OF TERM THREE, 2016

PHYSICS PAPER 1

2 hours 30 minutes.

INSTRUCTIONS TO CANDIDATES:

Attempt five questions, including at least one, but not more than two from each of the sections A, B, and C.

Assume where necessary:

Acceleration due to gravity, g = 9.81 ms^{-2} Electron charge, e = $1.6 \times 10^{-19} \text{ C}$

Stefan's Boltzmann's constant, σ = 5.7x10⁻⁸ Wm⁻² K⁻⁴

Radius of the earth $= 6.4 \times 10^6 \text{m}$ Radius of the sun $= 7.0 \times 10^8 \text{m}$ Radius of Earth's orbit about the sun $= 1.5 \times 10^{11} \text{m}$ Speed of light in a vacuum, $c = 3.0 \times 10^8 \text{ms}^{-1}$ Specific heat capacity of water $= 4,200 \text{J kg}^{-1} \text{ K}^{-1}$

Universal gravitational constant, G = $6.67 \times 10^{-11} \text{N m}^2 \text{ kg}^{-2}$ Avogadro's number N_A = $6.02 \times 10^{23} \text{ mol}^{-1}$

Density of water = 1000 kgm^{-3} Planck's constant, h = $6.63 \times 10^{-34} \text{ Js}$ Charge to mass ratio, e/m = $1.8 \times 10^{11} \text{ C kg}^{-1}$ The constant $1/4\pi\epsilon_0$ = $9.0 \times 10^9 \text{ F}^{-1}\text{m}$

Permittivity of free space, ϵ_0 = 8.85 x 10^{-12} Fm⁻¹ Mass of the electron = 9.11 x 10^{-31} kg

	SECTION A		
1(a)	What is meant by the following terms as applied to materials?		
	(i) tensile stress	(1)	
	(ii) tensile strain	(1)	
	(iii) elastic limit	(1)	
(b) (3)	Explain the occurrence of work hardening in an otherwise ductile	matei	rial.
(c)	Describe an experiment to determine Young's modulus for a steel wir	re. (5)	
	A rod of cross section area A and Young's modulus E is fixed in rigid Given that the linear expansivity of the rod is α derive an expression in it if the temperature changes by $\delta\theta$. (3)		orts at its e force
point	A metal wire of diameter 2.0×10^{-4} m and length 2 m has its ends find B 1.6 m apart, AB being horizontal. What force should be applied of the wire to pull the point of application to a point 0.65 m below lus = 2×10^{11} Nm ⁻²] (6)	ed at th	
2 (a)	(i) What are the distinguishing characteristics between <i>uniform circumotion</i> and <i>simple harmonic motion</i> ?	<i>lar</i> (3)	
radius	(ii) Derive an expression for the acceleration of a particle describing a s r with a uniform speed, v. (5)	circle	of
	A particle of mass m is hung on a string of length, l. It is then projected ontal velocity, V.	ed	with a
subse	(i) Show that when the string makes an angle θ with the vertical, duriquent motion, the tension in the string is	ng	the
	$m\{V^2/I - g(2 - 3cos\theta)\}$	(4)	
	(ii) Hence show that if the particle is to describe a complete circle, the	en	
	$V^2 > 5gI$	(3)	
(c)	The displacement x of a particle is given by $x = a \sin \omega t$		
	(i) Derive an expression for its velocity in terms of its displacement,		
	angular velocity, ω , and the amplitude, a .	(3)	
	(ii) Show that the total energy of the particle is constant.	(2)	

3(a)	(i) What is meant by acceleration due to gravity?	(1)	
(1)	(ii) Sketch the displacement-time graph for a body thrown vertically up	o in	air.
	(iii) What is meant by a trajectory?	(1)	
air at a	(iv) Show, from first principles, that the path followed by a particle angle, α , to the horizontal is parabolic. (3)	projec	ted in
(b)	A projectile is fired horizontally from the top of a cliff, 200 m high. The projectile lands 1000 m from the bottom of the cliff. Find the:	ne	
	(i) Initial speed of the projectile.	(4)	
	(ii) Velocity of the projectile just before it hits the ground.	(5)	
(c) helica	Describe an experiment to determine the acceleration due to gravity us l spring of unknown force constant. (5)	sing	a
4(a)	(i) Account for the existence of intermolecular forces.	(2)	
substa	(ii)Sketch a graph of potential energy against separation of two molecul nce and explain the main features of the graph. (4)	es	in a
(b)	(i) Define surface tension.	(1)	
	(ii)Show that the excess pressure, p, in an air bubble inside a liquid over outside pressure is given by	r	

2γ

p =

r

	where $ r $ is the radius of the bubble and $ \gamma $ its surface tension.	(3)
(c)	(i) State Archimedes principle.	(1)
of the	(ii) Show that the Up thrust on a body immersed in a fluid is equal to the fluid displaced. (4)	e weight
(d)	A solid weighs 237.5g in air and 12.5g when totally immersed in a liquid relative density 0.9. Calculate;	l of
	(i) the relative density of the solid	(2)
its volu	(ii) the relative density of a liquid in which the solid would float with or time exposed above the liquid surface. (3)	ne- fifth of
	SECTION B	
5(a)	Define <i>thermo- conductivity</i> of a material.	(1)
(b)	(i) Explain, using the molecular theory of matter the mechanism of heaconduction in insulators.	at (3)

insula	(ii) Briefly account for the fact that metals are better conductors ators. (3)	s of heat	than
(c) betwe	A wall of a building consists of two brick layers each of thicknessen which there is a layer of air 2.0 cm thick.	ess 10.0 cm	and
	Find the rate of heat flow through 1 m ² of the wall if the inner a temperatures of the building are 25°C and 15°C respectively.	and outer	
	[Thermal conductivities of brick and air are respectively, 0.7 W	m-1K-1 and	
	$0.024~{\rm Wm^{-1}K^{-1}}]$	(6)	
	(i) What is meant by a <i>black body</i> ? (ii) The average distance of Pluto from the sun is 40 times that out. If the sun radiates as a black body at 6000K, and is a calculate the surface temperature of Pluto. (6)	(1) of the earth 1.5×10^{11} m f	from From the
6(a)	(i) What is meant by cooling correction?	(1)	
heat o	(ii) Explain how the cooling correction may be estimated in the capacity of poor conductor of heat by the method of mixtures.	determinatio	n of the
(b)	(i) Define the term specific heat capacity of a substance.	(1)	

	(ii)An electrical heater rated 500W is immersed in a liquid of mass 2.0 I contained in a large thermos flask of heat capacity 840J K ⁻¹ at 28 ⁰ C. It is supplied to the heater for 10 minutes. If the specific heat capacity of	Electrical
_	13 J kg ⁻¹ K ⁻¹ , its specific latent heat of vaporization is 8.54 x 10 ³ J kg	-
	point is 78°C, estimate the amount of liquid which boils off stating any	•
made.	(7)	assumptions
made.		
(c)	(i) With reference to an electrical thermometer, describe the steps invol-	ved in
setting	up a Kelvin scale of temperature. (3)	
ı	(ii)The resistance of the element of a platinum resistance thermometer is	s 4.00Ω
	· ·	on the
	m resistance scale would correspond to a resistance of 9.84Ω ?	
1	(3)	
7(a)	Explain why	
	(i) at low pressures a real gas approximates to a perfect gas.	(2)
,	(1) at low pressures a real gas approximates to a perfect gas.	(2)
((ii) evaporation causes cooling.	(2)
		, ,
(b)	(i) State three properties of a saturated vapour.	(3)
	/**	. 1 1
	(ii) Air is trapped between a piston and a liquid in a cylinder. The piston	•
_	•	the liquid
	porated. Sketch a graph of pressure against volume for the content er and explain its features. (3)	its of the
cymnuc	t und explain to reacutes. (0)	

(iii) Describe an experiment to show that saturated vapour pressure vatemperature. (6)	ries with
(c) The saturated vapour pressure of ether at 0° C is 185 mmHg and at 20 440 mmHg. The bulb of a constant-volume gas thermometer contains dry a ether for saturation. If the observed pressure in the bulb is 1000 mmHg will it be at 0° C? (4)	ir and sufficient
SECTION C	
8(a) Define the following terms as used in photo electricity.	
(i) work function.	(1)
(ii) stopping potential	(1)
(b) (i)Describe an experiment to determine Planck's constant in a laborate	tory. (5)
(ii) Write down Einstein's photo-electric equation and explain how it emission of electrons from a metal surface. (3)	accounts for

	gnetic radiation of frequency 8 V. Calculate the maximum sp	-	
from the surface.	(4)	-	
(c)	P		
		_	
	C	-	
		1	Fig.1
Figure 1, shows two parallel metal plates P and Q each of length 5.0 cm and separated by a distance of 4.0 cm. A p.d. of 12V is applied between P and Q and the space between P and Q is a vacuum. A beam of electrons of speed $u = 1.0 \times 10^6 \text{ ms}^{-1}$ is directed midway between P and Q. Calculate the angle at which the beam emerges from the space between P and Q to the initial direction of the beam.			
(6)9(a) (i) Distinguish	ı between X-rays and cathode	rays.	(2)
(ii) Draw a we production of the X-ra	ll-labeled diagram of the X-ra ays.	y tube and outline the (6)	principles of
(b) (i) State Bragg	's law of X-ray diffraction.		(1)

set of p	(iii) A monochromatic beam of X-rays of wa parallel atomic planes of spacing $3.00 \times 10^{-10} \mathrm{m}$	•	
diffrac			
(c)	Briefly distinguish between X-ray line speci	tra and optical line spectra. (2)	
(d)	Figure 2 below shows some of the energy le	vels of a mercury atom	
	Energy in eV		
	0.0	- E _σ	
	-1.6		
	-3.7	_ E ₃	
	-5.5	- E ₂	
	-10.4	E ₁	
	Fig	g. 2	
	Determine		
(3)	(i) the minimum speed of an electron that co	ould just ionize the mercury	atom.
E2.	(ii) the wavelength of the radiation emitted	in an electron transition from (3)	E ₃ to

- 10(a) (i) Sketch and label the I-V characteristic for gaseous conduction. (2)
 - (ii) Explain the main features of the curve. (3)
- (b) A valve diode D was connected to rectify a.c supply to a load as shown in figure 3. C is a capacitor.




Fig. 3

- (i) State the purpose served by the capacitor C. (1)
- $\label{eq:continuous} \mbox{(ii) Explain the effect on the p.d's V_d and V_L of introducing a little } \qquad \mbox{mercury} \\ \mbox{vapour in the vacuum of D.} \qquad \qquad \mbox{(3)}$

(c)	A sinusoidal voltage of amplitude 0.1V is applied to the grid of triode of amplification factor 12. If the anode resistance of the triode is 15 k Ω , F	
	(i) Voltage will appear across a load of 18 $\mbox{k}\Omega$	(3)
	(ii) The gain of the stage at this load resistance.	(2)
(d)	(i) Draw a labeled diagram of a cathode ray oscilloscope (C.R.O)	(3)
	(ii) Describe how an alternating p.d may be measured using a C.R.O	(3)

BEGINNING-OF-TERM TEST

May 2012

S6 PHYSICS

Paper 1

2 ½ hours

NAME	Stream	H'se
Signature		

Answer FIVE questions in all, but not more than Three from one section.

Acceleration due to gravity, g = 9.81 ms^{-2}

Mass of the earth = $5.97 \times 10^{24} \text{ kg}$

Stefan's Boltzmann's constant, σ = 5.67x10⁻⁸ Wm⁻² K⁻⁴

Radius of the earth = 6.4×10^6 m

Radius of the sun = $7.0 \times 10^8 \text{m}$

Radius of Earth's orbit about the sun	=	$1.5 \text{ x} 10^{11} \text{m}$

Specific heat capacity of water = $4,200 \text{J kg}^{-1} \text{ K}^{-1}$

Universal gravitational constant, G = $6.67 \times 10^{-11} \text{N m}^{-2} \text{ kg}^{-2}$

Density of water = 1000 kgm^{-3}

Gas constant , R = $8.31 \text{ J mol}^{-1}\text{K}^{-1}$

Question	1	2	3	4	5	TOTAL
Marks						


SECTION A

1. (a) Define the following:		
	(i) Tensile stress	(1)	
	(ii) Tensile strain	(1)	
((iii) Young's modulus	(1)	
(b curve) (i) Sketch the stress-strain curve for a copper wire and explain the char . (4)	racteri	stic features of the
	(ii) How does the curve in b(i) compare with that of a material like glass	s?	(2)

- (c) Derive an expression for the energy stored in a unit volume of a stretched elastic material, expressing your answer in terms of Young's modulus and the strain of the material.(4)
- (d) A cylindrical copper rod of length 0.5 m and diameter 2 cm is fixed between two rigid supports at a temperature of 20°C. The temperature of the rod is raised to 70°C. Calculate

(i) the force exerted on the rigid supports, at this temperature.	(4)
(ii) the energy stored in the rod at 70°C	(3)
Young's modulus for copper = $1.2 \times 10^{11} \text{ Nm}^{-2}$ (Mean coefficient of line 20°C and 70°C is $1.7 \times 10^{-5} \text{ K}^{-1}$)	ear expansion of copper between
2. (a) Define simple harmonic motion.	(1)
(b) A particle moves so that its displacement x after time t is given by	у
$x = r \sin 2\pi t$	
(i) Show that the particle executes simple harmonic motion	(3)
(ii) Using the same axes sketch graphs of displacement vs time, v time. (3)	velocity vs time and acceleration vs
(c) A uniform cylinder of height h floats vertically in water with ¾ of if the cylinder is given a small vertical displacement and then released motion of period $\pi V\left(\frac{3h}{g}\right)$	•
	,
(d) A cube of mass 100 g rests on a light scale pan that is supported 24.5 Nm ⁻¹ . The pan is then set to oscillate vertically.	d on a vertical spring of constant
Find the maximum amplitude for the cube to remain in contact with the (3)	he pan throughout the motion.
(e) (i) Distinguish between free and damped oscillations and sketch each (4)	n a displacement-time graph of
3. (a) (i) State stokes' law of viscosity	(1)
(ii) A ball of density σ and radius r is released to fall through a liq Derive an expression for its terminal velocity during its motion. (3)	• •

- (iii) Two identical rain drops are observed to fall with a terminal velocity of 0.2 ms⁻¹ If the two coalesce into one drop, what will be the new terminal velocity? (4)
 - (b) (i) State Bernoulli's principle (2)
 - (ii) Explain one application of Bernoulli's principle (3)
 - (c) The diagram below shows a pipe delivering water



The cross-sectional area at section A is 30 cm² and the velocity there is 4 ms⁻¹ while the cross-sectional area at B is 15 cm². Find

- (i) the volume of water flowing per second (2)
- (ii) the static pressure at B if the atmospheric pressure is $1 \times 10^5 \text{ Nm}^{-2}$. (5)
- 4. (a) (i) Show that in uniform circular motion the accelerating force is towards the centre of the circle and derive an expression for the acceleration. (4)

(2)	(ii) State two differences between difficility circular motion and s	imple narmonic mode	JII.
	(ii) Explain briefly the action of a centrifuge.	(3)	
	(iii) A cyclist is to turn a corner of a horizontal circular track of racion between the tyres and the track surface if μ . For no skidding h dition that		
	$v < \sqrt{\mu rg}$	(3)	
(k	o) (i) Define gravitational potential.	(1)	
an o	(ii) Find an expression for the total energy of a satellite of mass rorbit of radius R.	m circling the earth of (3)	mass M in
(4)	(iii) Explain the effect of friction between such a satellite and the	atmosphere in which	it moves.
	SECTION B		
5. (a	a) Describe an experiment to verify Newton's law of cooling.	(5)	
(b	b) (i) Distinguish between <i>a real</i> and <i>an ideal</i> gas.	(3)	
	(ii) Derive the expression:		
P	= $\frac{1}{3}\rho \overline{c^2}$ for the pressure of an ideal gas of density ρ and mean	square speed $\overline{c^2}$. (6	5)
	c) (i) Explain why the pressure of a fixed mass of gas in a closed corperature of the container is raised.	ntainer increases whe	en
raise	(ii) Nitrogen gas is trapped in a container by a movable piston. If ed from 0° C to 50° C at constant pressure of 4.0 x 10^{5} Pa and the to	·	he gas is
3.0 >	x 10 ⁴ J, calculate the work done by the gas.		
[The	e molar heat capacity of nitrogen at constant pressure is 29.1 J mo	$I^{-1} K^{-1}; C_p/C_v = 1.4]$ (4	1)

6. (a) (i) What is meant by a <i>reversible process</i> ?	(1)
(ii) Explain the conditions for a reversible adiabatic change.	(3)
(iii) Explain why a gas heats up when it is compressed adiabatically	(3)
(b) The temperature of one mole of oxygen gas at a pressure of 3.0×10^{-2} when the gas expands adiabatically. Find the final pressure of the gas.	0 ⁵ Pa falls from 80°C to 17°C
[Take γ = 1.40]	(4)
(c) (i) Sketch a P-V curve for a real gas undergoing compression below it (1)	s critical temperature.
(ii) Explain the main features of the curve.	(3)
(d) When air saturated with water vapour in a rigid cylinder, at 100°C ar	
2.0×10^5 Pa pressure, is cooled to 20° C the pressure drops to 7.98×10^4 Pa the cylinder, calculate the saturated vapour pressure of water at 20° C. [x 10^5 Pa] (5)	
7. (a) State Prevost's theory of exchange with reference to heat radiation	. (1)
(b) Explain why metals are better conductors of heat compared to wo	od. (4)
(c) A is a composite slab made up of two layers of thickness d_1 and d_2 and k_2 respectively. B is a slab of thickness $d_1 + d_2$ with thermal conductivi slabs is maintained by heating at a temperature θ_1 and the other end at θ_2 unit area through the two slabs is the same, find k in terms k_1 , k_2 , d_1 and d_2 (6)	ty k. One face of each of the ₂ . If the rate of conduction per
(d) The heat radiation received by the earth from the sun is 1.4×10^3 Wt the sun emits as a black body, estimate the temperature of the sun. (4)	m ⁻² . Assuming this 90% of what

(e) With the aid of a labeled diagram, describe the construction pyrometer.	and mode of operation of the optical (5)		
SECTION C			
8. (a) (i) Sketch the I-V characteristic for gaseous conduction.	(2)		
(ii) Explain the main features of the characteristic.	(3)		
(b) (i) What is meant by thermionic emission?	(1)		
(ii) Describe an experiment to show that cathode rays conve	ey negative charge. (4)		
(c) A horizontal beam of electrons, moving with a uniform speed between two horizontal parallel charged plates.	ed, enters a uniform electric field		
Show that the path of the beam between the plates is a parabola	. (4)		
(d) In a Millikan's oil drop experiment a single negatively charged drop of radius 6×10^{-6} m was found to fall under gravity at a terminal velocity of 0.004 cm s ⁻¹ and to rise at 0.012 cm s ⁻¹ when a field of 2 x 10^{5} Vm ⁻¹ was suitably applied. Given that the viscosity of the medium was 2.122×10^{-5} Nsm ⁻² determine the number of electrons on the drop. (5)			
END			
1. (a) State Newton's laws of motion.	(3)		

- (b) Use Newton's laws of motion to show that linear momentum is conserved when two particles collide directly. (6)
- (c) Balls A, B and C of respective masses m_1 , m_2 and m_3 , lie in a straight line on a smooth surface. The balls are initially at rest. Ball A which is projected with a velocity V_1 towards B, makes an elastic collision with B. If B moves and makes a perfectly inelastic collision with C, show that both B and C move with a velocity

$$V_3 = \frac{2m_1m_2V_1}{(m_1 + m_2)(m_2 + m_3)}$$
(5)

- (d) The foot of a uniform ladder of weight 200 N is on a horizontal ground, and the top rests against a smooth vertical wall. A man of weight 800 N stands on the ladder one quarter of its length from the bottom. If the inclination of the ladder to the horizontal is 30° , find:
 - (i) the reaction at the wall (3)
 - (ii) the total force at the ground. (3)

P510/1 PHYSICS Paper 1

July, 2012

UGANDA ADVANCED CERTIFICATE OF EDUCATION MOCK EXAMINATIONS, 2012

PHYSICS

PAPER 1

2 hours 30 minutes.

INSTRUCTIONS TO CANDIDATES:

Attempt **five** questions, including at least **one**, but not more than **two** from each of the sections **A**, **B**, and **C**.

Assume where necessary.

Permittivity of free space ϵ_0 = 8.85 x 10^{-12} Fm⁻¹

Acceleration due to gravity, $g = 9.81 \text{ ms}^{-2}$

Electron charge, $e = 1.6 \times 10^{-19} \text{ C}$

Stefan's Boltzmann's constant, $\sigma = 5.7 \times 10^{-8} \text{ Wm}^{-2} \text{ K}^{-4}$

Radius of the earth $= 6.4 \times 10^6 \text{m}$

Radius of the sun = $7.0 \times 10^8 \text{m}$

Radius of Earth's orbit about the sun = $1.5 \times 10^{11} \text{m}$

Speed of light in a vacuum, $c = 3.0 \times 10^8 \text{ms}^{-1}$

Specific heat capacity of water = $4,200 \text{ J kg}^{-1} \text{ K}^{-1}$

Universal gravitational constant, G = $6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$

Avogadro's number N_A = $6.02 \times 10^{23} \text{ mol}^{-1}$

Density of water = 1000 kgm^{-3}

Gas constant, $R = 8.31 \text{ J mol}^{-1}\text{K}^{-1}$

Charge to mass ratio, e/m = 1.8 x 10¹¹ C kg⁻¹

The constant $\underline{1}$ = $9.0 \times 10^9 \, F^{-1} m$

4πεο


SECTION A

1. (a) (i) Define the coefficient of surface tension. (1)

(ii) Lycopodium powder is sprinkled over the surface of water in a dish. A small drop of soap solution is then introduced at the centre of the surface.

Explain what is observed on the surface. (3)

(b) Describe an experiment to investigate the effect of temperature on su tension of a liquid.	urface (6)
(c) (i) State Bernoulli's principle.	(1)
(ii) Explain how Bernoulli principle is applied in a filter pump.	(3)
(d) Water flows steadily along a uniform tube of cross-section 30 cm² . The pressure is 1.25 x 10 ⁵ Pa and the total pressure is 1.30 x 10 ⁵ Pa . Find	
(i) the flow velocity	(4)
(ii) the mass flow rate.	(2)
2. (a) Define the following:	
(i) A couple.	(1)
(ii) Torque.	(1)
(b) Show that when a couple rotates a body the work done is equal to the	ne product of the
torque and the angle in radians turned through.	(4)
(c) The diagram below shows three coplanar forces acting on a particle.	




2N

Find the magnitude, and inclination to the **12N** force, of an additional force that will bring about equilibrium.

(6)

(d) The figure below shows a uniform rigid plank AB of weight 100 N and length 5m freely hinged at end B in a vertical wall. A light rope is fixed between end A and a point C, 5 m vertically above B, so that AB makes an angle of 30° with the horizontal.



A box of weight **200 N** is hung from point **D** on the plank **2 m** from end **B**. Find;

(i) the tension in the rope.

(ii) the reaction at the hinge. (5)

3. (a) (i) Derive an expression for the tangential speed of a particle describing a circle of radius r, at an angular velocity ω . (3)

(3)

(ii) A coin is placed on a horizontal turntable at a point 5 cm from the centre of the turntable. If the coefficient of friction between the coin and the turntable is 0.5 , find the maximum frequency of the turntable for the coin to remain at the same point. (3)		
(b) (i) Define gravitational potential.	(1)	
(ii) Find an expression for the total energy of a satellite of mass ${\bf M}$ in an orbit of radius ${\bf R}$.	m circling the earth of (3)	
(iii) Explain the effect of friction between such a satellite and the it moves.	e atmosphere in which 4)	
(c) A particle executing simple harmonic motion vibrates in a straig the speeds of the particle are 4 ms ⁻¹ and 2 ms ⁻¹ when the particle is 3 c respectively from the equilibrium position, calculate the;		
(i) amplitude of oscillation.	(3)	
(ii) frequency of the particle.	(3)	

4. (a) (1) What is meant by a conservative force?	(1)
(ii) Show that if a body is moving in the gravitational field, the mechani is conserved. 4)	cal energ
(b) A particle is projected with a speed of 30 ms ⁻¹ to hit a target at a helabove the horizontal through the point of projection and at a distance of	ight of 12 m 60 m.
Find:	
(i) the angle of projection.	(5)
(ii) the time taken to hit the target.	(2)
(c) (i) What is meant by uniform acceleration?	(1)
(ii) A particle is projected vertically upwards from the top of a tree. Sketc	
displacement-time and a velocity-time graph for the motion up to the particle hits the ground.	(2)
(d) A car, which is slowing down steadily, passes a point A with a speed of	72 km
h -1 at the instant a motorcyclist at A is starting off in the same direction.	If the
motorcyclist maintains a steady acceleration of 2.5 ms ⁻² and catches up at point B when the car's speed is 18 km h ⁻¹ , find;	with the car
(i) the time taken to move from A to B .	(3)

(ii) the distance AB	(2)
SECTION B	
5 (a) (i) Define thermal conductivity.	(1)
(ii) Explain the mechanism of heat transfer in metals.	(3)
(iii) Describe an experiment to determine the thermal conductivity of a metal.	(6)
(b) (i) What is a black body?	(1)
(ii) Draw sketch graphs to show the variation of relative intensity of black bod radiation with wavelength for different temperatures.	y (2)
(iii) Explain the appearance of a metal ball placed in a dark room when its temp progressively raised from room temperature to just below melting. (3)	oerature is
(c) A heating element in form of a cylinder 30 cm long and 1.5 cm in diameter houtput of 1.8 kW . If its radiation is 85% that of a black body, find its temperature (4)	

	What assumptions are necessary in the derivation of tressure of an Ideal gas?	the Kinetic theory (4)	expression for
a wa	A beam of 2×10^{22} Nitrogen atoms, each of mass 2.32 all of a cubical container of edge 5.0 cm. The beam is an speed of the atoms is 500 ms^{-1} , find the pressure example (4)	reflected through 1	80 °. If the
(c) ((i) State Dalton's law of partial pressures.		(1)
(ii)Two containers A and B of volume 3×10^3 cm³ and 6×10^3 cm³ respectively contain helium gas at a pressure of 1.0×10^3 Pa and temperature 300 K. Container A is heated to 373 K while container B is cooled to 273 K. Find the final pressure of the Helium gas. (5)			
	Use the kinetic theory of gases to explain the effect of the control of the contr	of increasing tempe (3)	rature on
	Sketch a pressure versus volume curve for a real gas critical temperature. Explain the main features of th	0 0 1	
7.	(a) (i) Define specific latent heat of fusion.		(1)
	(ii) Describe an experiment to determine the specifi	ic latent heat of fus (5)	ion of water.

from that at constant volume.	(3)
(ii) Derive an expression for the difference of the above n	nolar heat capacities. (4)
(c) (i) State the conditions necessary for a reversible adiabate	atic process. (2)
(ii) A gas initially occupying a volume of 1.0 litre at a tempressure of 1.40 x 10 ⁵ Pa expands isothermally to a volum compressed adiabatically to its original volume. If $\gamma = 1.4$ f and pressure.	e of 2.0 litres . It is then
SECTION C	
8. (a) (i) What is meant by ionization energy of an atom?	(1)
(ii) Explain why ionization energy is always quoted as a nega	ative value. (2)
(b) Below are some of the energy levels of a mercury atom	
Energy in eV	
$E_{\infty} = 0.0$	

$E_4 = -1.6$		-	
$E_3 = -3.7$			
$E_2 = -5.5$		-	
П 10.4			
$E_1 = -10.4$			
Determine;			
(i) the <i>min</i>	imum speed of an electron th	at could just ionize the mercu	ry atom.(3)
(ii) the wav	velength of the radiation emit	ted in an electron transition fr	om E 3 to E 2.
		(3)	
	e principles of generation of o	continuous and line spectra of	
ray tube.		(5)	
	romatic beam of X-rays of want of a potassium chloride crystal	welength 2.0×10^{-10} m is incid	ent on a set of 3.1 x 10 ⁻¹⁰ m
-	cular mass of potassium chlori		0.1 x 10 III
(i) the glar	ncing angle for first order diff	raction maxima to be observed	l. (2)
(ii) the den	nsity of the potassium chloride	<u>a.</u>	(4)
()	,		(-)
0 () () 5 : :			(2)
9. (a) (1) Distingi	uish between <i>radioactivity an</i>	d nuclear fission.	(2)

(2)	
(b) Describe how a <i>Geiger-Muller tube</i> works.	(6)
(c) 1.2g of a substance form a point source of γ-rays. Only 1 in 10 ¹² of its radioactive and the half-life is 100 days . A Geiger-Muller tube facing the distance of 10 cm gives a count rate of 11 s ⁻¹ . If the window of the tube h cm ² find	e source at a
(i) the number of radioactive atoms present.	(5)
(ii) the mass number of the substance.	(2)
(d) Determine whether the nucleus ²¹⁰ Po is stable or it may undergo	
84	
disintegration to produce ^{206}Pb and an $\alpha\text{-particle}.$	(3)
82	
Mass of 210 Po = 209.937u	
Mass of 206 Pb = $205.929u$	
Mass of ${}^{4}\text{He} = 4.002u$	
10. (a) (i) Sketch the <i>I-V characteristic</i> for gaseous conduction.	(2)

(ii) Why are neutrons preferred to charged particles for inducing nuclear reactions.

(ii) Explain the main features of the curve.	(3)
(b) A beam of electrons moving with a velocity \mathbf{v} , enters normally a strong field, which is uniform and of induction \mathbf{B} . If each electron has a mass \mathbf{m} are assuming that the electrons suffer a deviation of 180° by the time they leave magnetic field,	nd charge e ,
(i) derive an expression for the distance d , covered by an electron while mofield. (3)	oving in the
(ii) write down the relationship between ${\bf d}$ and ${\bf B}$.	(1)
(c) (i) Draw a labeled diagram of a cathode ray oscilloscope (C.R.O)	(3)
(ii) Describe how an alternating p.d may be measured using a C.R.O	(3)
(d) In a Millikan's oil drop experiment a single negatively charged drop of ra	dius

END

viscosity of the medium was $2.122 \times 10^{-5} \, \text{Nsm}^{-2}$ determine the $\,$ number of electrons on the

rise at **0.012 cm s**⁻¹ when a field of **2 x 10** ⁵ **Vm**⁻¹ was suitably applied.

drop.

 $6 \ x \ 10^{-6} \ m$ was found to fall under gravity at a terminal velocity of $0.004 \ cm \ s^{-1}$ and to

5)

Given that the

P510/1 PHYSICS

Paper 1

UGANDA ADVANCED CERTIFICATE OF EXAMINATIONS

PHYSICS

PAPER 1 (PRACTICE PAPER)

2 hours 30 minutes.

INSTRUCTIONS TO CANDIDATES:

Attempt **five** questions, including at least **one**, but not more than **two** from each of the sections **A**, **B** and **C**.

Non-programmable scientific electronic calculators may be used.

Assume where necessary.

Permittivity of free space ε_0 = 8.85 x 10⁻¹² Fm⁻¹

Acceleration due to gravity, $g = 9.81 \text{ ms}^{-2}$

Electronic charge, e = 1..6 x10⁻¹⁹ C

Mass of the earth = $5.97 \times 10^{24} \text{ kg}$

Planck's constant, h = 6.6 x 10⁻³⁴ Js

Stefan's Boltzmann's constant, σ = 5.7x10⁻⁸ Wm⁻² K⁻⁴

Radius of the earth $= 6.4 \times 10^6 \text{m}$

Radius of the sun = $7.0 \times 10^8 \text{m}$

Radius of Earth's orbit about the sun = 1.5×10^{11} m

Speed of light in a vacuum, $c = 3.0 \times 10^8 \text{ms}^{-1}$

Thermal conductivity of aluminium = 210Wm⁻¹ K⁻¹

Specific heat capacity of water = 4,200J kg⁻¹ K⁻¹

Universal gravitational constant, G = $6.67 \times 10^{-11} \text{N m}^2 \text{ kg}^{-2}$

Avogadro's number N_A = 6.02 X 10²³ mol⁻¹

Density of water = 1000 kgm^{-3}

Gas constant, $R = 8.31 \text{ J mol}^{-1}\text{K}^{-1}$

Charge to mass ratio, e/m = 1.8 x 10¹¹ C kg⁻¹

The constant 1 = $9.0 \times 10^9 \,\text{F}^{-1}\text{m}$

 $\frac{}{4\pi\epsilon_0}$

SECTION A

1.	(a) Define the terms: <i>elastic limit</i> and <i>Young modulus.</i>	(2)
	(b) Explain the energy transformations that occur when a wire is stretched;	
	(i) elastically.	(3)
	(ii) plastically.	(2)
	(c) A rod of mild steel of uniform cross sectional area $0.0030\ m^2$ and length $1.0\ m$ stretched steadily until it breaks.	is
	(i) Sketch a graph to show the relationship between the force and the extension. Explain the shape of the graph.	(4)
	(ii) When the applied force on the rod is $120 kN$ the strain is $4.0 x 10^{-4}$. Calculate Young modulus for steel.	(3)
	(iii) The rod is found to break at a stretching force of $\bf 240~kN$. Explain why the stress at the section of the rod where the break occurs is like be much greater than $\bf 80000~kN~m^{-2}$.	ly to (2)
	(iv) Just before the rod breaks, its extension is about 4 cm . Estimate the work do in stretching the rod by this amount. Suggest why the calculated value of the word done is likely to be less than the actual value.	

2.	(a) State Kepler's laws of planetary motion.	(3)
	(b) Derive the expression for the centripetal force on mass ${\bf m}$ moving with unifor speed ${\bf v}$ in a circular path of radius ${\bf r}$.	rm (4)
	(c) Explain why:	
	(i) the earth has atmosphere.	(2)
	(ii) a cyclist has to lean over towards the centre of a bend in order to turn well.	(2)
	(d) A satellite of mass $120~kg$ moves in a circular orbit around the earth with a period of $7.7~x~10^{27}s$.	
	(i) Find its height above the earth.	(5)
	(ii) Calculate the mechanical energy of the satellite.	(4)
3.	(a) (i) Define surface tension and show that it is equal to the work done per unit area in increasing the liquid surface under isothermal conditions.	(3)
	(ii) Find the energy required to break up a drop of water of radius 0.5 cm into d of water each of radius 1 mm . [Surface tension of water is 2.27 x 10 ⁻² Nm ⁻¹]	rops (4)
	(b) Describe an experiment to measure surface tension of a liquid by the capillar	y
	tube method.	(6)

- (c) (i) Derive an expression for the excess pressure in a soap bubble of radius \mathbf{r} if the soap solution has surface tension $\mathbf{\gamma}$.
- (ii) The pressure inside an air bubble formed 40 cm below the water surface is $1.043 \times 10^5 \text{ Pa}$. If the surface tension of water is $2.27 \times 10^{-2} \text{ Nm}^{-1}$, find the radius of the bubble. [Atmospheric pressure = $1.013 \times 10^5 \text{ Pa}$] (4)
- 4. (a) What is meant by *simple harmonic motion*? (1)
 - (b) A cylindrical vessel of cross-sectional area **A** contains air of volume **V**, at pressure **P**, trapped by a frictionless piston of mass **M**. The piston is pushed down and then released.
 - (i) If the piston oscillates with simple harmonic motion, show that its frequency ${\bf f}$ is given by

$$f = \frac{A}{2\pi} \sqrt{\frac{P}{MV}} \tag{6}$$

- (ii) Show that the expression for f_i in b(i) is dimensionally correct. (2)
- (c) A mass of **0.1kg** suspended from a spring of force constant **24.5 Nm**-1 is pulled vertically downwards through a distance of **5.0cm** and released. Find the

(ii) position of the mass **0.3 seconds** after release. (4)

(d) Sketch the following graphs for a body performing simple harmonic motion:	
(i) Velocity against displacement	(1)
(ii) Displacement against time.	(1)
(e) With the aid of sketch graphs, distinguish between free and damped oscillation	ns. (3)
SECTION B	
(a) (i)Explain why the difference between the principal heat capacities of a solid negligible.	is (2)
(ii) Derive an expression for the difference in molar heat capacities of a gas.	(4)
(b) (i)Explain why the sound propagation in air is considered to be an adiabatic process.	(2)
(ii) A fixed mass of gas at a pressure P_1 and volume V_1 expands isothermally to a pressure P_2 and volume V_2 . Derive an expression for the work done by the gas.	(4)
(c) A gas of volume 2 litres at a temperature of 27°C and pressure of 1.5 x 10⁵ F heated at constant pressure until its volume doubles. It is then cooled at constant volume back to its original temperature before finally being compressed isothermally to its original volume.	
Draw a p-V diagram of the whole cycle and find the net work done by the gas.	(5)


5.


- (d) The pressure **P** of an ideal gas of density ρ , is given by $P = \frac{1}{3} \rho \overline{c^2}$, where $\overline{c^2}$ is the mean-square speed of its molecules. Use this expression to derive Avogadro's hypothesis. (3)
- 6. (a) State Prevost's theory of exchange with reference to heat radiation. (1)
 - (b) Explain why metals are better conductors of heat compared to wood. (3)
 - (c) A is a composite slab made up of two layers of thickness d_1 and d_2 with thermal conductivities k_1 and k_2 respectively. B is a slab of thickness $d_1 + d_2$ with thermal conductivity k. One face of each of the slabs is maintained by heating at a temperature θ_1 and the other end at θ_2 . If the rate of conduction per unit area through the two slabs is the same, find k in terms k_1 , k_2 , d_1 and d_2 . (6)
 - (d) The heat radiation received by the earth from the sun is $1.4 \times 10^3 \, Wm^{-2}$. Assuming this 90% of what the sun emits as a black body, estimate the temperature of the sun.
 - (e)(i) With the aid of a labeled diagram, describe the construction and mode of operation of the optical pyrometer. (5)
 - (ii) State how the range of an optical pyrometer can be extended. (1)
- 7. (a) (i) The specific heat capacity of copper is **400J** kg⁻¹ K⁻¹. Explain the meaning of the statement. (1)

	(ii) State the factors that determine the rate of cooling of a body.	(2)
	(iii) Explain why a small body cools faster than a bigger one of the same material and shape. (4)	l
minim (7)	(b) With the aid of a well labeled diagram, describe an electrical method of determi the specific latent heat of vaporization of water, stating the precautions taken to ize heat losses	ning
5°C.	(c) Two taps, one delivering water at a temperature of 25° C at a rate of 0.3 g s^{-1} and other delivering water at a temperature of 15° C at a rate of 0.4 g s^{-1} , are directed to well-lagged copper calorimeter of mass 50 g containing ice at 0° C while stirring. After minutes the contents of the calorimeter are found to have attained a temperature of Find the mass of ice originally in the calorimeter. Specific latent heat of fusion of ice = $3.36 \times 10^5 \text{ Jkg}^{-1}$ (6)	a r 5
	SECTION C	
8. (a	a) (i) Sketch the I-V characteristic for gaseous conduction and explain the main features of the curve.	(4)
((ii) Explain why this is not a good method for producing cathode rays.	(2)
	(b) (i) What is meant by thermionic emission ?	(1)
	(ii) Describe an experiment to show that cathode rays travel in straight lines.	(4)

- (c) A horizontal beam of electrons, moving with a uniform speed, enters a uniform electric field between two horizontal parallel charged plates. Show that the path of the beam between the plates is a parabola. (4)
- (d) A charged oil drop of mass $6 \times 10^{-15} \, kg$ falls vertically in air with a steady velocity between two parallel vertical plates $5 \, mm$ apart. When a p.d of $3000 \, V$ is applied between the plates, the drop now falls with a steady velocity at an angle 58.5° to the vertical. Calculate the charge Q on the drop. (Neglect the up thrust on it)

- 9. (a) State the characteristics of photoelectric emission. (4)
 - (b) The figure below shows graphs of current I against p.d V applied across a photoelectric cell that receives monochromatic light for two different intensities A and B (A < B)





10.

- (i) Draw a diagram to show the set-up you could use to establish these results. (3) (ii) State what V_T represents and explain the features of the graphs. (3) (c) Describe an experiment to determine Planck's constant in a laboratory. (5) (d) Light of frequency **5.0** x **10**¹⁴ Hz liberates electrons with maximum energy 2.31×10^{-19} J from a certain metallic surface. What is the wavelength of ultra-violet light which liberates electrons of maximum energy **8.93** x **10**⁻¹⁹ J from the surface? (5) (a) (i) Distinguish between **radioactivity** and **nuclear fission**. (2) (ii) Define **binding energy** of a nucleus (1) (b) (i) What is half life of a radioactive substance? (1)
 - (ii) Derive the relationship between half life and the decay constant of a radioactive substance. (3)
 - (c) With the aid of a diagram describe how a Wilson type cloud chamber detects ionizing particles. (6)

- (d) A nucleus of uranium ${\bf 238}$ disintegrates with the emission of a nucleus ${\bf X}$ and an alpha particle.
- (i) Write down the equation for the reaction. (1)
- (ii) Find the velocity with which the alpha particle is released, assuming the uranium nucleus was at rest before disintegration.

Mass of 238 U = 238.12492 u

Mass of X = 234.11650 u

Mass of ${}^{4}\text{He} = 4.00387$

$$1 \text{ u} = 1.66 \text{ x } 10^{-27} \text{ kg} = 1.494 \text{ x } 10^{-12} \text{ J}$$
 (6)

END

P510/1 PHYSICS Paper 1

July, 2012

UGANDA ADVANCED CERTIFICATE OF EDUCATION MOCK TWO (2), 2012

PHYSICS

PAPER 1

2 hours 30 minutes.

INSTRUCTIONS TO CANDIDATES:

Attempt **five** questions, including at least **one**, but not more than **two** from each of the sections **A**, **B**, and **C**.

Assume where necessary:

Permittivity of free space ϵ_o = 8.85 x 10^{-12} Fm⁻¹

Acceleration due to gravity, $g = 9.81 \text{ ms}^{-2}$

Electron charge, $e = 1.6 \times 10^{-19} \text{ C}$

Stefan's Boltzmann's constant, $\sigma = 5.7x10^{-8} Wm^{-2} K^{-4}$

Radius of the earth = 6.4×10^6 m

Radius of the sun = $7.0 \times 10^8 \text{m}$

Radius of Earth's orbit about the sun = 1.5×10^{11} m

Speed of light in a vacuum, $c = 3.0 \times 10^8 \text{ms}^{-1}$

Specific heat capacity of water = $4,200 \text{J kg}^{-1} \text{ K}^{-1}$

Universal gravitational constant, G = $6.67 \times 10^{-11} \text{N m}^{-2} \text{ kg}^{-2}$

Avogadro's number N_A = $6.02 \times 10^{23} \text{ mol}^{-1}$

Density of water = 1000 kgm^{-3}

Planck's constant, h = $6.63 \times 10^{-34} \text{ Js}$

Charge to mass ratio, e/m = 1.8 x 10¹¹ C kg⁻¹

The constant . 1 = $9.0 \times 10^9 \text{ F}^{-1}\text{m}$


4πεο

SECTION A

- 1. (a) (i) State the principle of conservation of linear momentum. (1)
- (ii) Show how Newton's laws of motion may be used to arrive at the principle of conservation of momentum. (5)
- (b) The engine of a 5-tonne lorry can develop 30kW, and its maximum speed on level road is 90 kmh⁻¹. Assuming that the friction resistance is constant, calculate the greatest speed at which the lorry can climb a hill of 1 in 25. (4)

(c) Explain the principle of rocket propulsion. (3)

(d) The figure below shows a block A of mass 1 kg resting on a horizontal plane. Another block B of mass 2 kg on a plane inclined at an angle of 30° to the horizontal is connected to A by a string passing over a smooth pulley P. All the parts of the string are parallel to the respective planes.



Given that the coefficient of friction at each point of contact is 0.3, find

(ii) A ball of density σ and radius r is released to fall through a liquid of density ρ and viscosity η . Derive an expression for its terminal velocity during its motion. (3)

(iii) Describe a simple experiment to demonstrate stream line and turbu in a liquid.	llent flow (4)
(b) (ii) Given that the volume of a liquid issuing per second from a pipe do on the co-efficient of viscosity η , the radius of the pipe a and the pressure grashow that	-
Volume per second $= \underline{kpa^4}$	
ηl	(5)
 (c) (i) Given that the surface tension of a soap solution is 2.7 x 10⁻² Nm⁻¹, the work done in forming 10 soap bubbles each of diameter 1.6 cm. (ii) Two soap bubbles, one of radius x and the other of larger radius y, g attached to each other. 	(3)
Find an expression for the radius of the common interface in terms of x and y	y. (4)
3. (a) (i) State Newton's law of gravitation.	(1)
(ii) Deduce the dimensions of the Universal gravitational constant.	(2)

(b) Assume that a planet moves with uniform speed in a circular orbit of radius r about the sun. The periodic time of the planet is T. Show that the quantity (r^3/T^2) is independent of the mass of the planet. (4)

(c) An artificial satellite is launched in a circular orbit round the equator at a height of 3.60×10^7 m above the surface of the Earth.		
(i) Determine from first principles the speed with which the satellite must launched to maintain it in the orbit.	t be (5)	
(ii) Determine the periodic time of the satellite.	(3)	
(iii) What deduction can be made from the result obtained in (ii) above wit reference to the satellite?	th (2)	
(iv) If another satellite moves in a circular orbit concentric with that of the above satellite, at a height of 1.48×10^7 m above the surface of the Earth, what is to periodic time of this second satellite?		
4. (a)(i) Account for the existence of intermolecular forces.	2)	
(ii)Sketch a graph of potential energy against separation of two molecules in a	l	
substance and explain the main features of the graph. (4)		
(c) A hydrometer floats in pure water with a volume V submerged. When in a liquid of density ρ , it floats with the liquid level at a distance X above the water mark. If the stem has a uniform cross sectional area ϕ , derive an expression for X in terms of V, ρ , and ϕ . (4)		

(b) Derive an expression for energy stored per unit volume of a stretched elastic material, expressing your answer in terms of young's modulus and the strain of the material. (4)

(c) A cylindrical copper rod of length 0.5m and of diameter 4.0×10^{-2} m is fixed between two rigid supports at a temperature of 20.0° C. The temperature of the raised to 70.0° C		
(i) Calculate the force exerted on the rigid supports at this temperature	(4)	
(ii) What is the energy stored in the rod at 70°C?	(3)	
Young's modulus for copper = $1.2 \times 10^{11} \text{Nm}^{-2}$		
Mean coefficient of linear expansion of copper between 20°C and 70°C =1.7 x	<i>10⁻⁵K⁻¹</i>	
SECTION B		
5. (a) (i) Mention the steps involved in establishing a temperature scale. (2)		
(ii) State the advantages and disadvantages a thermocouple thermometer a platinum resistance one.	has over (3)	
(iii) Describe how the temperature of a liquid bath may be measured usin platinum resistance thermometer.	g a (4)	
(b) (i) The resistance R_θ of platinum varies with temperature $\theta^o C$ as measure constant-volume gas thermometer according to the equation $R_o(1+8000\alpha\theta-\alpha\theta^2) \text{ where } \alpha \text{ is a constant. Calculate the temperature on the}$	ed by a R _θ =	
platinum scale corresponding to 400°C on this gas scale. (4)		
(ii) Explain why the two thermometers do not agree exactly. (1)		

(c) In a continuous flow experiment on a liquid of specific heat capacity 4200 Jkg⁻¹K⁻¹ the following results were obtained:

Ammeter reading/A	Voltmeter reading/V	Mass of liquid collected per min/g
2.00	25.2	75.0
2.52	30.0	115.9

If the inflow temperature was 15°C find

- (i) the outflow temperature. (4)
- (ii) the rate of heat loss. (2)
- 6. (a) (i) What is meant by cooling correction? (1)
 - (ii) Explain how the cooling correction may be estimated in the determination of the heat capacity of poor conductor of heat by the method of mixtures. (5)
 - (b) (i) Define the term specific heat capacity of a substance. (1)
 - (ii)An electrical heater rated 500W is immersed in a liquid of mass 2.0kg contained in a large thermos flask of heat capacity 840J $\rm K^{-1}$ at 28°C. Electrical power is supplied to the heater for 10 minutes. If the specific heat capacity of the liquid is $\rm 2.5~x10^3Jkg^{-1}K^{-1}$, its specific latent heat of vaporization is $\rm 8.54~x$

boils off stating any assumptions made. (7)
(c) With the aid of a diagram, describe how a total radiation pyrometer can be used to measure the temperature of a very hot body. (6)
7. (a) (i) State the conditions for an adiabatic change.(2)
(ii) Explain why a gas cools when it undergoes an adiabatic expansion. (2)
(iii) Show that an adiabatic is steeper than an isothermal on the same P-V diagram. (3)
(b) Describe an experiment to verify Charles's law.(6)
(c) Derive the <i>ideal gas equation</i> . (4)
(d) A rigid vessel contains air initially at atmospheric pressure. It is then inflated using a pump whose barrel has an effective volume equal to a $\frac{1}{20}$ th of the vessel's volume. Find the number of stroke to be performed before the pressure triples. (3)
8. (a) Define the following terms as used in photo-electricity.

 $10^{3} J\ kg^{\text{--}1}$ and its boiling point is $78^{\circ} C,$ estimate the amount of liquid which

(i) work function. (1 mark)

(ii) **stopping potential** (1 mark)

- (b) Using the quantum theory, explain the laws of photo-electric emission. (3)
 - (c) (i)Describe an experiment to determine Planck's constant. (5)

(ii)Electromagnetic radiation of frequency 8.8 x 10¹⁴ Hz falls upon a surface whose work function is 2.5eV. Calculate the maximum kinetic energy of photoelectrons released from the surface. (4)

(c)

Two parallel plates P and Q each of length 4.0cm and separated by a distance of 4.0cm in vacuum, have p.d. of 12V applied between them. A beam of electrons of speed $u = 1.0 \times 10^6 \text{ ms}^{-1}$ is directed midway between P and Q. Find the angle at which the beam emerges from the space between P and Q to the initial direction of the beam.

- 9. (a) (i) What is meant by the term 'binding energy of a nucleus'? (1)
- (ii) Calculate the binding energy per nucleon of an α particle, expressing your result in MeV.

Mass of proton = 1.0080u

Mass of a neutron = 1.0087u

Mass of an α - particle = 4.0026u

1u = 931 MeV. (4)

(iii) Sketch a graph of binding energy per nucleon against mait to explain liberation of energy by nuclear fusion and nuclear fiss (6)	
(b) Derive an expression relating the half life of a radioactive mat decay constant λ .	terial, T½ and the
(c) When ²³⁸ ₉₂ U decays, the end product is ²⁰⁶ ₈₂ Pb. The half life is 1.4	4×10^{17} s.
Suppose a rock sample contains ²⁰⁶ ₈₂ Pb and ²³⁸ ₉₂ U in the ratio 1: 5 by the	weight. Calculate
(i) number of ²⁰⁶ ₈₂ Pb atoms in 5.0 g of the rock sample.	(3)
(ii) age of the rock.	(4)
10. (a) A high p.d is applied across two electrodes in air container tube. Describe, with the aid of labeled diagrams, what will be observessure in the tube is progressively reduced down to very low pressively.	rved when the
(b) List four main properties of cathode rays.	(4)
(c) A charged oil drop of mass $3.27 \times 10^{-15} \text{kg}$ is held stationary horizontal metal plates across which a p.d of 1000 V is applied. If t plates is 1.5 cm find the number of electrons on the drop. (5)	

(d) With the aid of a labeled diagram describe the principle of operation of an ionisation chamber. (6)

UGANDA ADVANCED CERTIFICATE OF EDUCATION MOCK TWO (2), 2012

PHYSICS PAPER 1

2 hours 30 minutes.

INSTRUCTIONS TO CANDIDATES:

Attempt **five** questions, including at least **one**, but not more than **two** from each of the sections **A**, **B**, and **C**.

Assume where necessary:

Permittivity of free space ϵ_0 = 8.85 x 10^{-12} Fm⁻¹

Acceleration due to gravity, g = 9.81 ms⁻²

Electron charge, $e = 1.6 \times 10^{-19} \text{ C}$

Stefan's Boltzmann's constant, $\sigma = 5.7x10^{-8} Wm^{-2} K^{-4}$

Radius of the earth = $6.4 \times 10^6 m$

Radius of the sun = $7.0 \times 10^8 \text{m}$

Radius of Earth's orbit about the sun = $1.5 \times 10^{11} \text{m}$

Speed of light in a vacuum, $c = 3.0 \times 10^8 \text{ms}^{-1}$

Specific heat capacity of water = $4,200 \text{J kg}^{-1} \text{ K}^{-1}$

Universal gravitational constant, G = $6.67 \times 10^{-11} \text{N m}^{-2} \text{ kg}^{-2}$

Avogadro's number N_A = $6.02 \times 10^{23} \text{ mol}^{-1}$

Density of water = 1000 kgm^{-3}

Planck's constant, h = $6.63 \times 10^{-34} \text{ Js}$

Charge to mass ratio, e/m = 1.8 x 10¹¹ C kg⁻¹

The constant $\frac{1}{4\pi\varepsilon_0}$ = 9.0 x 10⁹ F⁻¹m


SECTION A

- 1. (a) (i) State the principle of conservation of linear momentum. (1)
 - (ii) Show how Newton's laws of motion may be used to arrive at the principle of conservation of linear momentum. (5)
 - (b) The engine of a **5-tonne** lorry can develop **30kW**, and its maximum speed on level road is **90 kmh**⁻¹. Assuming that the frictional resistance is constant, calculate the greatest speed at which the lorry can climb a hill of **1** in 25.

(4)

(c) Explain the principle of rocket propulsion. (3)

(d) The figure below shows a block A of mass 1 kg resting on a horizontal plane. Another block B of mass 2 kg on a plane inclined at an angle of 30° to the horizontal is connected to A by a string passing over a smooth pulley P. All the parts of the string are parallel to the respective planes.



Given that the coefficient of friction at each point of contact is 0.3, find

(i) the acceleration of the blocks.

(5)

 $\mbox{(ii)}$ the tension in the string .

(2)

2. (a) (i) Define *coefficient of viscosity* of a liquid.

- (ii) A ball of density σ and radius r is released to fall through a liquid of density ρ and viscosity η . Derive an expression for its terminal velocity during its motion. (3)
- (iii) Describe a simple experiment to demonstrate stream line and turbulent flow in a liquid. (4)
- (b) (ii) Given that the volume of a liquid issuing per second from a pipe depends on the co-efficient of viscosity η , the radius of the pipe a, and the pressure gradient, show that;

Volume per second $= \underline{kpa^4}$

$$\eta l$$
 (5)

(c) (i) Given that the surface tension of a soap solution is $2.7 \times 10^{-2} \, Nm^{-1}$, calculate the work done in forming 10 soap bubbles each of diameter $1.6 \, cm$.

(3)

- (ii) Two soap bubbles, one of radius x and the other of larger radius y, get attached to each other. Find an expression for the radius of the common interface in terms of x and y. (4)
- 3. (a) (i) State Newton's law of gravitation. (1)

(ii) Deduce the dimensions of the Universal gravitational constant. (2)
(b) Assume that a planet moves with uniform speed in a circular orbit of radius rabout the sun. The periodic time of the planet is T . Show that the quantity (r³/T²) is independent of the mass of the planet. (4)
(c) An artificial satellite is launched in a circular orbit round the equator at a height of 3.60×10^7 m above the surface of the Earth.
(i) Determine from first principles the speed with which the satellite must be launched to maintain it in the orbit. (5)
(ii) Determine the periodic time of the satellite. (3)
(iii) What deduction can be made from the result obtained in (ii) above with reference to the satellite? (2)
(iv) If another satellite moves in a circular orbit concentric with that of the above satellite, at a height of 1.48×10^7 m above the surface of the Earth, what is the periodic time of this second satellite? (3)
4. (a)(i) Account for the existence of intermolecular forces. (2)

- (ii)Sketch a graph of potential energy against separation of two molecules in a substance and explain the main features of the graph. (3)
- (b)Describe an experiment to determine Young Modulus for a steel wire. (4)
- (c) Derive an expression for energy stored per unit volume of a stretched elastic material, expressing your answer in terms of Young modulus and the strain of the material. (4)
- (d) A cylindrical copper rod of length **0.5m** and of diameter **4.0 x 10⁻² m** is fixed between two rigid supports at a temperature of **20.0°C**. The temperature of the rod is raised to **70.0°C**
 - (i) Calculate the force exerted on the rigid supports at this temperature.

(4)

(3)

(ii) What is the energy stored in the rod at **70°C**?

Young modulus for copper = $1.2 \times 10^{11} \text{Nm}^{-2}$

Mean coefficient of linear expansion of copper between 20 $^{\circ}$ C and 70 $^{\circ}$ C =1.7 $\times 10^{-5}$ K⁻¹

SECTION B

5. (a) (i) Mention the steps involved in establishing a temperature scale. (2)

- (ii) State the advantages and disadvantages a thermocouple thermometer has over a platinum resistance one. (3)
- (iii) Describe how the temperature of a liquid bath may be measured using a platinum resistance thermometer. (4)
- (b) (i) The resistance R_θ of platinum varies with temperature θ °C as measured by a constant-volume gas thermometer according to the equation $R_\theta = R_o(1 + 8000\alpha\theta \alpha\theta^2)$ where α is a constant. Calculate the temperature on the platinum scale corresponding to 400 °C on this gas scale. (4)
 - (ii) Explain why the two thermometers do not agree exactly. (1)
- (c) In a continuous flow experiment on a liquid of specific heat capacity **4200** J **kg**⁻¹ K⁻¹ the following results were obtained:

Ammeter	Voltmeter	Mass of liquid collected	
reading (A)	reading (V)	per min (g)	
2.00	25.2	75.0	
2.52	30.0	115.9	

If the inflow temperature was 15 °C, find;

(i) the outflow temperature.	(4)
(ii) the rate of heat loss.	(2)
6. (a) (i) What is meant by cooling correction ?	(1)
(ii) Explain how the cooling correction may be estimated in the determine the heat capacity of poor conductor of heat by the method of mixture (5)	
(b) (i) Define the term specific heat capacity of a substance.	(1)
(ii)An electrical heater rated 500W is immersed in a liquid of macontained in a large thermos flask of heat capacity 840 J K ⁻¹ at 28 power is supplied to the heater for 10 minutes. If the specific heat the liquid is 2.5 x 10 ³ J kg ⁻¹ K ⁻¹ , its specific latent heat of vaporizat 10 ³ J kg ⁻¹ and its boiling point is 78 °C, estimate the amount of liquid boils off stating any assumptions made.	°C. Electrical at capacity of tion is 8.54 x
(c) With the aid of a diagram, describe how a total radiation pyrometer used to measure the temperature of a very hot body. (6)	er can be

7. (a) (i) State the conditions for an adiabatic change.	(2)
(ii) Explain why a gas cools when it undergoes an adiaba (2	_
(iii) Show that an adiabatic is steeper than an isothermal diagram.	on the same P-V (3)
(b) Describe an experiment to verify Charles's law.	(6)
(c) Derive the <i>ideal gas equation</i> .	(4)
(d) A rigid vessel contains air initially at atmospheric pressurinflated using a pump whose barrel has an effective volute vessel's volume. Find the number of stroke to be perfortriples. (3)	ume equal to a $\frac{1}{20}$ th of
SECTION C	
8. (a) Define the following terms as used in photo-electricity.	
(i) work function.	(1)

(ii) stopping potential.	(1)
(b) Using the quantum theory, explain the	laws of photo-electric emission. (3)
(c) (i)Describe an experiment to determine	Planck's constant. (5)
(ii)Electromagnetic radiation of freque whose work function is 2.5eV . Calculate photoelectrons released from the surfac	e the maximum kinetic energy of
(d)Two parallel plates P and Q each of leng distance of 4.0cm in vacuum, have p beam of electrons of speed u = 1.0 x 10 6 ms Find the angle at which the beam emerges the initial direction of the beam. (6)	.d. of 12V applied between them. A -1 is directed midway between P and Q .
9. (a) (i) What is meant by the term 'bindi	ng energy of a nucleus? (1)
(ii) Calculate the binding energy per your result in MeV .	nucleon of an α particle, expressing
Mass of proton	= 1.0080u
Mass of a neutron	= 1.0087u

Mass of an $lpha$ - particle	= 4.0026u	
1u	= 931 MeV.	(4)
(iii) Sketch a graph of binding energy use it to explain liberation of energy by nucl		number and ear fission.
(b) Derive an expression relating the half l the decay constant, λ .	ife of a radioactive materi (6)	ial, T½ and
(c) When $^{238}_{92}$ U decays, the end product is 2	$^{06}_{82}$ Pb. The half life is 1.4 x	10 ¹⁷ s.
Suppose a rock sample contains $^{206}_{82}$ Pl weight. Calculate the:	and $^{238}_{92}$ U in the ratio 1: 5	5 by
(i) number of $^{206}_{82}$ Pb atoms in 5.0 g o	f the rock sample.	(3)
(ii) age of the rock.		(4)
10. (a) A high p.d is applied across two ele	ectrodes in air contained i	n a closed glass

tube. Describe, with the aid of labeled diagrams, what will be observed when the

pressure in the tube is progressively reduced down to very low pressures.

(b) List **four** main properties of cathode rays.

(5)

(4)

(c) A charged oil drop of mass $3.27 \times 10^{-15} \text{kg}$ is held stationary between two horizontal metal plates across which a p.d of 1000 V is applied. If the separation of the plates is 1.5 cm, find the number of electrons on the drop.

(5)

(d) With the aid of a labeled diagram, describe the principle of operation of an ionization chamber. (6)

END