#### GRAPHS OF RATIONAL FUNCTIONS

A rational function is a fraction of polynomials of the type  $y = \frac{g(x)}{h(x)}$  where h(x) is strictly a

function of x. e.g 
$$y = \frac{5}{x+3}$$
,  $y = \frac{x+1}{2x+6}$ ,  $y = \frac{(x-1)(x-3)}{(x+1)(x-2)}$ ,  $y = \frac{x^2+4x+3}{x+2}$  e.t.c

## **Basic Investigations**

These provide information from which graphs are developed. These include;

1. Points of intersection of the curve and the axes.( x and y- intercepts).

## (a) x- intercepts,

The curve cuts the x- axis when y = 0

i.e. for the curve 
$$y = \frac{g(x)}{h(x)}$$
,  $\frac{g(x)}{h(x)} = 0$ 

$$\rightarrow g(x) = 0$$
 i.e. The numerator = 0.

## Example.

Find the x –intercepts of the following curves where applicable.

(i) 
$$y = \frac{5}{x+3}$$
 (ii)  $y = \frac{x+1}{2x+6}$  (iii)  $y = \frac{(x-1)(x-3)}{(x+1)(x-2)}$  (iv)  $y = \frac{x^2+4x+3}{x+2}$  (v)  $y = \frac{x^2+3}{x-1}$ 

#### Solution;

Note; (we shall just equate the numerators to zero where applicable)

i. Since the numerator  $\neq 0$  i.e equals 5 then, the curve has no x- intercepts.

ii. 
$$x + 1 = 0$$

$$\rightarrow x = -1$$
 Therefore the x –intercept is  $(-1,0)$ 

iii. 
$$(x-1)(x-3) = 0$$

$$\rightarrow x = 1, x = 3$$
 Therefore the x –intercepts are (1,0) and (3,0)

iv. 
$$x^2 + 4x + 3 = 0$$

$$(x+1)(x+3) = 0$$

$$\rightarrow x = -1, x = -3$$
 Therefore the x –intercepts are  $(-1,0)$  and  $(-3,0)$ 

v. Since  $x^2 + 3 = 0$  has **no real roots** i.e  $x^2 = -3$  then the curve has no x –intercepts.

# (b) y-intercepts,

The curve cuts the y- axis when x = 0

i.e. for the curve  $y = \frac{g(x)}{h(x)}$ , the y-intercept occurs at  $y = \frac{g(0)}{h(0)}$ . Implying to get

the y-ordinate we just substitute x = 0 in the expression  $y = \frac{g(x)}{h(x)}$ .

$$\rightarrow$$
 the y-intercept is the point  $\left(0, \frac{g(0)}{h(0)}\right)$ 

# Example.

Find the y –intercepts of the following curves where applicable.

(i) 
$$y = \frac{5}{x+3}$$
 (ii)  $y = \frac{x+1}{x(2x+1)}$  (iii)  $y = \frac{(x-1)(x-3)}{(x+1)(x-2)}$ 

## Solution:

**Note**; (we shall just substitute x = 0 in the expression given where applicable)

- i.  $y = \frac{5}{0+3} = \frac{5}{3}$ 
  - $\rightarrow$  the y-intercept is the point  $(0,\frac{5}{2})$
- $y = \frac{0+1}{0(2(0)+1)}$ . Here y is undefined since the **denominator equals zero** implying **no** y-intercept.

iii. 
$$y = \frac{(0-1)(0-3)}{(0+1)(0-2)} = \frac{-3}{2} = -1.5$$

- $\rightarrow$  the y-intercept is the point (0, -1.5)
- 2. Turning points of a curve.

These are points at which  $\frac{dy}{dx} = 0$ .

## **Examples**

Find the turning points of the curves below where applicable

(i) 
$$y = \frac{3(x-3)}{(x+1)(x-2)}$$
 (ii)  $y = \frac{x+1}{2x+6}$ 

(ii) 
$$y = \frac{x+1}{2x+6}$$

## Solution

i. For 
$$y = \frac{3(x-3)}{(x+1)(x-2)} = \frac{3x-9}{x^2-x-2}$$
,

For turning points, 
$$\frac{dy}{dx} = 0$$
.  

$$\frac{dy}{dx} = \frac{(x^2 - x - 2) \cdot 3 - (3x - 9) \cdot (2x - 1)}{(x^2 - x - 2)^2}$$

$$\frac{3x^2 - 3x - 6 - 6x^2 + 21x - 9}{(x^2 - x - 2)^2} = 0$$
$$-3x^2 + 18x - 15 = 0$$

$$-3x^2 + 18x - 15 = 0$$

$$x^{2} - 6x + 5 = 0$$
$$(x - 1)(x - 5) = 0$$

$$x = 1$$
 ,  $x = 5$ 

When 
$$x = 1$$
,  $y = \frac{3(1-3)}{(1+1)(1-2)} = 3$ 

When 
$$x = 5$$
,  $y = \frac{3(5-3)}{(5+1)(5-2)} = \frac{1}{3}$ 

Therefore the turning points are (1,3) and  $(5,\frac{1}{3})$ 

We shall continue straight to establish the nature of the turning points above As follows;

$$\frac{dy}{dx} = \frac{-3x^2 + 18x - 15}{(x^2 - x - 2)^2}$$

$$\frac{x}{\text{Sign of } \frac{dy}{dx}} - \frac{1}{0} + \frac{1}{0} + \frac{1}{0} - \frac{1}{0}$$
Min

... The turning points are  $(1,3)_{min}$  and  $\left(5,\frac{1}{3}\right)_{max}$ .

**Note:** the expression,  $y = \frac{3(x-3)}{(x+1)(x-2)}$  above could have also been differentiated by introducing natural logarithms as follows.

$$ln(y) = ln\left(\frac{3(x-3)}{(x+1)(x-2)}\right)$$
  

$$ln(y) = ln(3x-9) - ln(x+1) - ln(x-2)$$
  
But  $\frac{d}{dx}[lnf(x)] = \frac{f'(x)}{f(x)}$ ,

So we apply that logarithmic differentiation on all the terms in our expression above as follows

$$\frac{1}{y}\frac{dy}{dx} = \frac{3}{3x-9} - \frac{1}{x+1} - \frac{1}{x-2}$$

$$\frac{1}{y}\frac{dy}{dx} = \frac{3(x+1)(x-2) - (3x-9)(x-2) - (3x-9)(x+1)}{(3x-9)(x+1)(x-2)}$$

$$\frac{dy}{dx} = \frac{(-3x^2 + 18x - 15)}{(3x-9)(x+1)(x-2)} \times \frac{(3x-9)}{(x+1)(x-2)}$$

$$\frac{dy}{dx} = \frac{-3x^2 + 18x - 15}{[(x+1)(x-2)]^2}$$

$$\frac{dy}{dx} = \frac{-3x^2 + 18x - 15}{(x^2 - x - 2)^2}$$

ii. For 
$$y = \frac{x+1}{2x+6}$$

$$\frac{dy}{dx} = \frac{(2x+6).1 - (x+1).2}{(2x+6)^2} = 0$$

$$= \frac{4}{(2x+6)^2} = 0$$

$$4 = 0. (2x+6)^2$$

Implying there is no solution. Therefore the curve has no turning points.

#### 3. Restricted a rea.

This is the region on the graph where the curve cannot exist.

#### **Examples**

Find the restricted region for the curve given by  $y = \frac{3x-9}{x^2-x-2}$ 

#### **Solution:**

$$y = \frac{3x - 9}{x^2 - x - 2}$$

$$yx^2 - x(y+3) + 9 - 2y = 0$$

For non-real roots of x,

$$[-(y+3)]^2 - 4y(9-2y) < 0$$

$$9y^2 - 30y + 9 < 0$$

$$3y^2 - 10y + 3 < 0$$

$$(3y-1)(y-3) < 0$$

	$y < \frac{1}{3}$	$\frac{1}{3} < y < 3$	<i>y</i> > 3	
(3y - 1)	_	+	+	
(y - 3)	_	_	+	
(3y-1)(y-3)	+	_	+	

$$\therefore \quad \frac{1}{3} < y < 3$$

 $\therefore$  The function cannot lie between  $\frac{1}{3}$  and 3

## 4. Asymptotes

An asymptote is a line to which the curve tends to approach (but actually never touches it) as one or both of the x or y values tend(s) to infinity. Asymptotes can be;

- Vertical
- Horizontal
- Oblique (slanting)
- Curved asymptote.

# (a) Vertical asymptote.

A vertical asymptote is a vertical line to which the curve tends to approach (but actually never touches it) as y values tend to infinity.

To obtain vertical asymptotes, we equate the denominator of the rational function to zero and solve for *x* values.

## **Examples**;

Find the vertical asymptotes for the following functions where applicable;

(i) 
$$y = \frac{5}{x+3}$$

(ii) 
$$y = \frac{x+1}{x^2-4}$$


$$y = \frac{5}{x+3}$$
 (ii)  $y = \frac{x+1}{x^2-4}$  (iii)  $y = \frac{(x-1)(x-3)}{x^2-x-2}$ 

(iv) 
$$y = \frac{2}{x^2 - 3}$$
 (v)  $y = \frac{x^2 - x - 2}{x + 1}$ 

#### Solution.

We shall equate the denominator to zero.

- (i) x + 3 = 0x = -3
  - $\therefore$  The line x = -3 is the vertical asymptote.
- (ii)  $x^2 4 = 0$  $x^2 = 4$  $x = \pm 2$ 
  - $\therefore$  The lines x = -2 and x = 2 are the vertical asymptotes
- (iii)  $x^2 x 2 = 0$  (x+1)(x-2) = 0 x = -1, x = 2
  - $\therefore$  The lines x = -1 and x = 2 are the vertical asymptotes.
- (iv)  $x^{2} 3 = 0$  $x^{2} = 3$  $x = \pm \sqrt{3}$ 
  - $\therefore$  The lines  $x = -\sqrt{3}$  and  $x = \sqrt{3}$  are the vertical asymptotes.
- (v)  $y = \frac{x^2 x 2}{x + 1}$  $y = \frac{(x + 1)(x 2)}{(x + 1)}$ 
  - y = x 2, which is a line. But the original function is not defined at x = -1 so the function is aline not defined at a point (-1, -3)
  - .. The function has no vertical asymptote. As seen in the sketch below



#### (b) Horizontal asymptote.

A horizontal asymptote is a horizontal line to which the curve tends to approach as x values tend to  $-\infty$  or  $+\infty$  or both.

#### How to find the horizontal asymptotes for rational functions

• If the degree of the polynomial function of the numerator is less than the degree of the polynomial function of the denominator, then y = 0 (x - axis) is the horizontal asymptote of the rational function.

**Note;** The degree of a polynomial is the highest power of the variable (say x or y etc.) in the individual terms of the polynomial.

## Example.

Find the horizontal asymptote of the function  $y = \frac{x+1}{(x-1)(x+2)}$ 

#### Solution.

We must expand first to create polynomials in the numerator and denominator to get,  $y = \frac{x+1}{x^2+x-2}$ 

The degree of the numerator is 1 and that of the denominator is 2

Therefore the horizontal asymptote is the line y = 0 (x - axis).

If the degree of the polynomial function of the numerator is equal to the degree of the polynomial function of the denominator, then the horizontal asymptote of the rational function is;

 $y = \frac{\text{coefficient of the term with the highest power in the numerator}}{\text{coefficient of the term with the highest power in the denominator}}.$ 

#### **Example**

Find the horizontal asymptote of the function  $y = \frac{(6x-1)(x-3)}{(3x+1)(x-2)}$ 

#### Solution;

The function above expands to  $y = \frac{6x^2 - 19x + 3}{3x^2 - 5x - 2}$ 

The term with the highest power in the numerator is  $6x^2$  with coefficient 6 while the term with the highest power in the denominator is  $3x^2$  with coefficient 3.

The horizontal asymptote becomes  $y = \frac{6}{3} = 2$ 

Therefore the horizontal asymptote is the line y = 2.

**Note;** if the degree of the numerator polynomial is greater than the degree of the denominator polynomial, then the rational function does not have a horizontal asymptote but rather has either an oblique asymptote or curved asymptote as explained below.

#### (c) Oblique (slanting) asymptote.

An oblique asymptote is a slanting line to which the curve tends to approach as x and y values tend to  $-\infty$  or  $+\infty$ .

If the degree of the polynomial function of the numerator is greater than the degree of the polynomial function of the denominator by **1**, then the function has an oblique asymptote.

To obtain an oblique asymptote of the rational function, we first express the function in the form  $y = Q(x) + \frac{R(x)}{D(x)}$  using long division.

The oblique asymptote is the line y = Q(x).

#### Example;

Find the oblique asymptote of the curve given by  $y = \frac{x^2 + 4x + 3}{x + 2}$ .

#### Solution;

Clearly the degree of the numerator function is 2 and that of the denominator is 1 which gives a difference in degrees of 1 implying we shall have an oblique asymptote.

We shall first split the function  $y = \frac{x^2+4x+3}{x-1}$  using long division;

$$\begin{array}{c|c}
x+5 \\
x-1 & x^2+4x+3 \\
\underline{x^2-x} & 5x+3 \\
\underline{5x-5} & 8
\end{array}$$

This implies that  $y = x + 5 + \frac{8}{x-1}$ 

Therefore the oblique asymptote is the line y = x + 5

# (d) Curved asymptote;

A curved asymptote is a curve to which the function tends to approach as x and y values tend to  $-\infty$  or  $+\infty$ .

If the degree of the polynomial function of the numerator is greater than the degree of the polynomial function of the denominator by **more than 1**, then the function has a curved asymptote.

To obtain a curved asymptote of the rational function, we first express the function in the form  $y = Q(x) + \frac{R(x)}{D(x)}$  using long division.

The curved asymptote is the curve y = Q(x).

#### Example;

Find the curved asymptote of the rational function  $y = \frac{x^3+2}{x}$ .

#### Solution;

Clearly the degree of the numerator function is 3 and that of the denominator is 1 which gives a difference in degrees of 2 implying we shall have a curved asymptote.

We shall first split the function  $y = \frac{x^3+2}{x}$  using long division

$$\begin{array}{c|cccc}
x^2 \\
\hline
x & x^3 & + 2 \\
\hline
x^3 & & 2
\end{array}$$

This implies that  $y = x^2 + \frac{2}{x}$ 

Therefore the curved asymptote is the curve  $y = x^2$ 

# (e) The behaviour (sign) of the curve $y = \frac{g(x)}{h(x)}$ in the interval $-\infty < x < \infty$ .

To determine the sign of the function throught its domain, we construct a table.

The function, *y* can only change sign after the points where it cuts the x- axis and vertical asymptotes.

We shall use the points above to create regions of the curve in our table.

# Examples;

Establish the nature of the function  $y = \frac{(x+1)(x-6)}{(x+3)(x-2)}$ . Indicating clearly its sign in the various intervals of x.

## Solution;

$$x - \text{intercepts},$$
  
 $(x + 1)(x - 6) = 0$ 

$$x = -1, x = 6$$

Vertical asymptotes

$$(x+3)(x-2)=0$$

$$x = -3, x = 2$$

Region table.

	x < -3	-3 < x < -1	-1 < x < 2	2 < x < 6	<i>x</i> > 6
(x + 1)	_		+	+	+
(x - 6)	_		1	1	+
(x + 3)	_	+	+	+	+
(x-2)	_	_	_	+	+
(x+1)(x-6)	+	_	+	_	+
$y = \frac{1}{(x+3)(x-2)}$					

# Note;

- In the intervals where y is +, it implies that the curve is above the x axis
- In the intervals where y is -, it implies that the curve is below the x axis